



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07H	A2	(11) International Publication Number: WO 98/34942 (43) International Publication Date: 13 August 1998 (13.08.98)
(21) International Application Number: PCT/US98/02702 (22) International Filing Date: 6 February 1998 (06.02.98) (30) Priority Data: 08/797,472 6 February 1997 (06.02.97) US 08/798,508 10 February 1997 (10.02.97) US (63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Applications US 08/797,472 (CIP) Filed on 6 February 1997 (06.02.97) US 08/798,508 (CIP) Filed on 10 February 1997 (10.02.97) (71) Applicant (for all designated States except US): INSPIRE PHARMACEUTICALS, INC. [US/US]; 4222 Emperor Boulevard #470, Durham, NC 27703 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): PENDERGAST, William [US/US]; 5815 Williamsburg Way, Durham, NC 27713 (US). YERXA, Benjamin, R. [US/US]; 1330 Hathaway Road, Raleigh, NC 27608 (US). RIDEOUT, Janet, L. [GB/US]; 3101 Morningside Drive, Raleigh, NC 27607		(US). SIDDIQI, Suhaib, M. [AT/US]; 1005 Paddock Drive, Raleigh, NC 27609 (US). (74) Agent: HALLUIN, Albert, P.; Howrey & Simon, 1299 Pennsylvania Avenue, N.W., Box 34, Washington, DC 20004-2402 (US). (81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: CERTAIN DINUCLEOTIDES AND THEIR USE AS MODULATORS OF MUCOCILIARY CLEARANCE AND CILIARY BEAT FREQUENCY (57) Abstract <p>The present invention relates to certain novel dinucleotides and formulations thereof which are highly selective agonists of the P2Y₂ and/or P2Y₄ purinergic receptor. They are useful in the treatment of chronic obstructive pulmonary diseases such as chronic bronchitis, PCD, cystic fibrosis, as well as prevention of pneumonia due to immobility. Furthermore, because of their general ability to clear retained mucus secretions and stimulate ciliary beat frequency, the compounds of the present invention are also useful in the treatment of sinusitis, otitis media and nasolacrimal duct obstruction. They are also useful for treatment of dry eye disease and retinal detachment.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon	KR	Republic of Korea	PL	Poland		
CN	China	KZ	Kazakhstan	PT	Portugal		
CU	Cuba	LC	Saint Lucia	RO	Romania		
CZ	Czech Republic	LI	Liechtenstein	RU	Russian Federation		
DE	Germany	LK	Sri Lanka	SD	Sudan		
DK	Denmark	LR	Liberia	SE	Sweden		
EE	Estonia			SG	Singapore		

CERTAIN DINUCLEOTIDES AND THEIR USE AS MODULATORS OF MUCOCILIARY CLEARANCE AND CILIARY BEAT FREQUENCY

5

This application is a continuation-in-part application of U.S. Serial No. 08/798,508 filed February 10, 1997.

10

INTRODUCTION

Technical Field

This invention relates to certain dinucleotides which increase the hydration of retained mucus secretions, stimulate the production of mucins and increase ciliary beat frequency to increase clearance of retained secretions.

Background of the Invention

Chronic obstructive pulmonary disease (COPD) affects 15 million patients in the U.S. and is the sixth leading cause of death. It is characterized by the retention of mucus secretions in the lungs. Many patients diagnosed with COPD have a disorder called chronic bronchitis (CB), and 600,000 patients are hospitalized each year due to an acute exacerbation of CB. Cystic fibrosis and Primary Ciliary Dyskinesia (PCD) are other examples of lung disorders which assume a clinical profile similar to COPD. Ciliary dyskinesia, whether primary or secondary, results in retained secretions that can only be cleared by coughing.

Another disease state characterized by the accumulation of retained mucous secretions is sinusitis. Sinusitis is an inflammation of the paranasal sinuses typically associated with an upper respiratory infection. It is this country's most common health-care complaint, affecting an estimated 31 million people. (A. Moss and V. Parsons, National Center for Health Statistics, 1986: 66-7, DHHS Publication No. (PHS)86-1588 (1985)).

Otitis media (OM) is a viral or bacterial infection of the middle ear which primarily afflicts children under the age of three. It is usually precipitated by an upper respiratory infection which spreads into the middle ear via the nasopharynx and eustachian

tube. Approximately 25-50 million office visits are made each year for diagnosis and treatment of OM. By age three, about 75% of children will have had at least one episode of acute OM (J. Klein, *Clin. Infect. Dis.* 19, 823-33 (1994)). Following appropriate treatment with antibiotics, accumulated fluid in the middle ear remains, causing hearing impairment and potential language and cognitive development delays. Enhanced ability to clear secretions in the middle ear would reduce or eliminate significant sequelae of otitis media.

An additional disorder resulting from retained secretions is pneumonia. Patients who are immobilized for a variety of reasons are at high risk for developing pneumonia. Despite extra vigilance and numerous interventions, pneumonia develops in over 400,000 patients per year, with significant morbidity and mortality.

There are also situations where it is therapeutically desirable to increase drainage of the lacrimal system. When the lacrimal drainage system is not functioning properly the result can be excessive tearing (epiphora), mucopurulent discharge, and recurrent dacryocystitis. Current treatments for nasolacrimal duct obstruction are mostly invasive surgical procedures, and researchers have sought to discover noninvasive pharmaceutical treatments.

Tear secretion can be stimulated from lacrimal accessory tissues via P2Y₂ and /or P2Y₄ purinergic receptor-mediated mechanisms similar to those which hydrate airway epithelia. Dry eye disease is the general term for indications produced by abnormalities of the precorneal tear film characterized by a decrease in tear production or an increase in tear film evaporation, together with the ocular surface disease that results. Currently, the pharmaceutical treatment of dry eye disease is mostly limited to administration of artificial tears (saline solution) to temporarily rehydrate the eyes. However, relief is short lived and frequent dosing is necessary.

Normally, mucous secretions are removed via the mucociliary clearance (MCC) system. MCC relies on the integrated action of three components: 1) mucus secretion by goblet cells and submucosal glands; 2) the movement of cilia on epithelial cells

which propels the mucus across the luminal surface; and 3) ion transport into and out of luminal epithelial cells which concomitantly controls the flow of water into the mucus.

It is now known that nucleoside phosphates such as uridine 5'-triphosphate (UTP) modulate all of the components of the MCC system. First, UTP has been shown to increase both the rate and total amount of mucin secretion by goblet cells *in vitro* (M. Lethem, et al., *Am J. Respir. Cell Mol. Biol.* 9, 315-22 (1993)). Second, UTP has been shown to increase cilia beat frequency in human airway epithelial cells *in vitro* (D. Drutz, et al., *Drug Dev. Res.* 37(3), 185 (1996)). And third, UTP has been shown to increase Cl⁻ secretion, and hence, water secretion from airway epithelial cells *in vitro* (S. Mason, et al., *Br. J. Pharmacol.* 103, 1649-56 (1991)). In addition, it is thought that the release of surfactant from Type II alveolar cells in response to UTP (Gobran, *Am. J. Physiol.* 267, L625-L633 (1994)) contributes to optimal functioning of the lungs and may assist in maximizing MCC (M. Knowles, et al., *N. Engl. J. Med.* 325, 533-38 (1991)). UTP has been shown to increase intracellular Ca⁺⁺ due to stimulation of phospholipase C by the P2Y₂ receptor (H. Brown, et al., *Mol. Pharmacol.* 40, 648-55 (1991)).

UTP's modulation of all components of the mucociliary escalator system results in a 2.5-fold improvement in lung mucociliary clearance in normal volunteers without any significant side-effects (K. Olivier, et al., *Am J. Respir. Crit. Care Med.* 154, 217-23 (1996)). In addition, UTP significantly enhanced cough clearance (clearance of retained secretions by coughing) in patients with PCD (P. Noone, et al., *Am. J. Respir. Crit. Care Med.* 153, A530 (1996)).

Because of UTP's demonstrated ability to increase the clearance of retained mucous secretions, applicants were motivated to investigate whether other nucleoside phosphates could be equally, if not more, therapeutically effective. The present invention is based upon this investigation.

Previously described dinucleotides are listed in Table I, along with their corresponding literature references.

TABLE I

DINUCLEOTIDES IN THE LITERATURE

(numbers in parentheses correspond to references that follow)

Np ₂ N	Np ₂ N'	Np ₂ N	Np ₂ N'	Np ₂ N	Np ₂ N'
Ap ₂ A (4,1)	Ap ₂ NAD (6)	Up ₂ U (1)	Ap ₂ T (20)	Up ₂ U (2,3)	Ap ₂ U (3)
Gp ₂ G (5,1)	Ap ₂ TAD (6)	Ap ₂ A(1,4,29)	m ⁷ Gp ₂ G (5)	Ap ₂ A(1,4,29)	Ap ₂ C (3)
m ⁷ Gp ₂ m ⁷ G(5)	Ap ₂ C-NAD(6)	Xp ₂ X (1)	m ^{2,7} Gp ₂ G(5)	Cp ₂ C (3)	Ap ₂ G (3)
	Ap ₂ C-PAD(6)	m ⁷ Gp ₂ m ⁷ G(5)	m ^{2,7} Gp ₂ G(5)	Gp ₂ G (1,5)	Gp ₂ U (3)
	Ap ₂ BAD (6)	Gp ₂ G (1)		Xp ₂ X (1)	Gp ₂ C (3)
	m ⁷ Gp ₂ G (5)			Dp ₂ D (15)	Up ₂ C (3)
	Up ₂ U (43)			eAp ₂ eA (7)	Ap ₂ T (20)
				m ⁷ Gp ₂ m ⁷ G(5)	m ⁷ Gp ₂ G (5)
					m ^{2,7} Gp ₂ G (5)
					m ^{2,2,7} Gp ₂ G(5)

Np ₂ N	Np ₂ N'	Np ₂ N	Np ₂ N'	Np ₂ N
Ap ₂ A (4)	Ap ₂ T (20)	Ap ₂ A (4)	Ap ₂ T (20)	Ap ₂ A (4)

AppZppA	DppZppD	ApZppZpA	ApSpZpSpA
Z	Z	Z	Z
CH ₂ (8)	CH ₂ (15)	CH ₂ (8)	CHF (8)
CH ₂ CH ₂ (8)	CH ₂ CH ₂ (15)	CH ₂ CH ₂ (8)	CF ₂ (8)
CHF (8)	CHF (15)	CHF (8)	O (8)
CF ₂ (8)	CF ₂ (15)	CF ₂ (8)	
CHCl (8)	CHCl (15)	CHCl (8)	
CCl (8)	CCl (15)	CCl (8)	

15 A = Adenosine

U = Uridine

G = Guanosine

T = Thymidine

X = Xanthosine

TAD = Tiazofurin

20 BAD = Benzamide riboside

D = 2,6-Diaminopurine

eA = Ethenoadenosine

m⁷G = 7-Methylguanosinem^{2,7}G = 2,7-Dimethylguanosinem^{2,2,7}G = 2,2,7-Trimethylguanosine

NAD = nicotinamide riboside

C-NAD = C-nicotinamide riboside

C-PAD = C-picolinamide riboside

N = Nucleoside

25 (1) M.A.G. Sillero et al., Eur. J. Biochem., 76, 331 (1977)

(2) C.G. Vallejo et al., Biochim. Biophys. Acta, 483, 304 (1976)

- (3) H. Coste et al., J. Biol. Chem., 262, 12096 (1987)
- (4) K.E. Ng et al., Nucleic Acid Res., 15, 3573 (1987)
- (5) J. Stepinski et al., Nucleosides & Nucleotides, 14, 717 (1995)
- (6) A. Zatorski et al., J. Med. Chem., 39, 2422 (1996)
- 5 (7) P. Rotilan et al., FEBS, 280, 371 (1991)
- (8) P.C. Zamecnik et al., Proc. Natl. Acad. Sci., 89, 2370 (1992)
- (9) J. Walker et al., Biochemistry, 32, 14009 (1993)
- (10) R.H. Hideman et al., J. Biol. Chem., 266, 6915 (1991)
- (11) J. Luthje et al., Eur. J. Biochem., 173, 241 (1988)
- 10 (12) R.H. Silverman et al., Microbiological Rev., 43, 27 (1979)
- (13) C.D. Lobaton et al., Eur. J. Biochem., 50, 495 (1975)
- (14) G. Lowe et al., Nucleosides & Nucleotides, 10, 181 (1991)
- (15) G.M. Blackburn et al., Nucleosides & Nucleotides, 10, 549 (1991)
- (16) J.C. Baker et al., Mutation Res., 208, 87 (1988)
- 15 (17) G. Klein et al., Biochemistry, 27, 1897 (1988)
- (18) E. Castro et al., Br. J. Pharmacol., 100, 360 (1990)
- (19) D.R. Elmaleh et al., Proc. Natl. Acad. Sci., 81, 918 (1984)
- (20) R. Bone et al., J. Biol. Chem., 261, 16410 (1986)
- (21) Fed. Amer. Soc. Exper. Bio., Abstr. Part I, no. 1878 (1991)
- 20 (22) M.T. Miras-Portugal et al., Ann. NY Acad. Sci., 603, 523 (1990)
- (23) A. Guranowski et al., Biochemistry, 27, 2959 (1988)
- (24) F. Grummt et al., Plant Mol. Bio., 2, 41 (1983)
- (25) A.G. McLennan et al., Nucleic Acid Res., 12, 1609 (1984)
- (26) P. Zamecnik et al., Analytical Biochem., 134, 1 (1983)
- 25 (27) E. Rapaport et al., Proc. Natl. Acad. Sci., 78, 838 (1981)
- (28) T. Kimura et al., Biol. Pharm. Bull., 18, 1556 (1995)
- (29) E. Schulze-Lohoff et al., Hypertension, 26, 899 (1995)
- (30) B.K. Kim et al., Proc. Natl. Acad. Sci., 89, 11056 (1992)
- (31) P.C. Zamecnik et al., Proc. Natl. Acad. Sci., 89, 2370 (1992)
- 30 (32) H. Morii et al., Eur. J. Biochem., 205, 979 (1992)
- (33) E. Castro et al., Pflugers Arch., 426, 524 (1994)
- (34) H. Schluter et al., Nature, 367, 186 (1994)
- (35) E. Castro et al., Br. J. Pharmacol., 206, 833 (1992)
- (36) T. Casillas et al., Biochemistry, 32, 14203 (1993)
- 35 (37) J. Pintor et al., J. Neurochem., 64, 670 (1995)
- (38) E. Castro et al., J. Biol. Chem., 270, 5098 (1995)
- (39) V.A. Panchenko et al., Neuroscience, 70, 353 (1996)
- (40) E. Castro et al., Br. J. Pharmacol., 100, 360 (1990)
- (41) J. Pintor et al., Gen. Pharmac., 26, 229 (1995)
- 40 (42) J. Pintor et al., Br. J. Pharmacol., 115, 895 (1995)
- (43) A. Kanavarioti et al., Tett. Lett., 32, 6065 (1991)


45

SUMMARY OF THE INVENTION

5

The invention provides novel compounds of Formula I and pharmaceutical compositions thereof. The invention also provides compounds useful in the clearance of retained mucous secretion and the enhancement of ciliary beat frequency. Accordingly, a broad embodiment of the invention is directed to compounds of general Formula I or the pharmaceutically acceptable esters or salts thereof:

Formula I



wherein:

X is oxygen, methylene, difluoromethylene, imido;

$n = 0, 1$ or 2 ;

$m = 0, 1$ or 2 ;

$n + m = 0, 1, 2, 3$ or 4 ; and

B and B' are each independently a purine residue or a pyrimidine residue linked through the 9- or 1- position, respectively;

Z = OH or N_3 ;

Z' = OH or N_3 ;

Y = H or OH;

Y' = H or OH;


provided that when Z is N₃, Y is H or when Z' is N₃, Y' is H; and further provided that the compounds of Table I are excluded.

5 The compounds of the present invention are highly selective agonists of the P2Y₂ and /or P2Y₄ purinergic receptor; thus, they may be useful in the treatment of chronic obstructive pulmonary diseases such as chronic bronchitis, PCD, and cystic fibrosis, and may also be useful in the treatment of immobilized patients who are at risk for developing pneumonia. Furthermore, because of their general ability to clear retained mucus secretions and stimulate ciliary beat frequency, the compounds of the present invention may also be
10 useful in the treatment of sinusitis, otitis media and nasolacrimal duct obstruction. They may also be useful for the treatment of dry eye, retinal detachment and wound healing. In addition, because of the pharmacological actions of these compounds, they are useful in facilitating sputum induction procedures. Additionally, it is postulated that the compounds of
15 the present inventions could enhance the performance of athletes by increasing the clearance of mucous secretions from the lungs.

DETAILED DESCRIPTION OF THE INVENTION

This invention provides novel compounds of Formula I and pharmaceutical
20 compositions thereof. The invention also provides compounds useful in the clearance of retained mucous secretion and the enhancement of ciliary beat frequency. Accordingly, a broad embodiment of the invention is directed to novel compounds of general Formula I:

Formula I



25

wherein:

X is oxygen, methylene, difluoromethylene, imido;

n = 0, 1 or 2;

m = 0, 1 or 2;

n + m = 0, 1, 2, 3 or 4; and

B and B' are each independently a purine residue or a pyrimidine residue linked through the 9- or 1- position, respectively;

Z = OH or N₃;

Z' = OH or N₃;

Y = H or OH;

Y' = H or OH;


provided that when Z is N₃, Y is H or when Z' is N₃, Y' is H; and further provided that the compounds of Table I are excluded; or pharmaceutically acceptable esters or salts thereof.

The furanose sugar is preferably in the β-configuration.

The furanose sugar is most preferably in the β-D-configuration.

Preferred compounds of Formula I are the compounds of Formula IA

Formula IA



wherein:

- 5 X=O;
 n+m=1 or 2;
 Z, Z', Y and Y'=OH;
 B and B' are uracil, thymine, cytosine, guanine, adenine, xanthine, hypoxanthine or as
 defined in Formulas II and III; or
- 10 X=O;
 n+m=3 or 4;
 Z, Z', Y and Y'=OH;
 B=uracil;
15 B' is uracil, thymine, cytosine, guanine, adenine, xanthine, hypoxanthine or as defined
 in Formulas II and III; or
- X=O;
 n+m=1 or 2;
20 Z, Y and Y'=OH;
 Z'=H;
 B=uracil;
 B' is uracil, thymine, cytosine, guanine, adenine, xanthine, hypoxanthine or as
 defined in Formulas II and III; or
- 25 X=O;
 n+m=0, 1 or 2;
 Z and Y=OH;
 Z'=N₃;
30 Y'=H;
 B=uracil;
 B'=thymine; or
- X=O;
 n+m=0, 1 or 2;
35 Z and Z'=N₃;
 Y and Y'=H;
 B and B'=thymine; or
- 40 X=CH₂, CF₂ or NH;
 n and m=1;
 Z, Z', Y and Y'=OH;
 B and B' are is uracil, thymine, cytosine, guanine, adenine, xanthine, hypoxanthine or
 as defined in Formulas II and III;

45


provided that the compounds of Table I are excluded; or pharmaceutically acceptable salts

thereof.

- 5 Another preferred group of the compounds of Formula I are the compounds of Formula IB or the pharmaceutically acceptable salts thereof:

Formula IB

10



wherein:

X is oxygen, methylene, difluoromethylene or imido;

15 $n = 0$ or 1 ;


$m = 0$ or 1 ;

$n + m = 0, 1$ or 2 ; and

B and B' are each independently a purine residue, as in Formula II, or a pyrimidine residue, as in Formula III, linked through the 9- or 1- position, respectively. In the instance where B and B' are uracil, attached at N-1 position to the ribosyl moiety, then the total of $m + n$ may equal 3 or 4 when X is oxygen (see example 5). The ribosyl moieties are in the D- configuration, as shown, but may be L-, or D- and L-. The D- configuration is preferred.

25

Formula II



wherein:

R_1 is an alkyl or aryl moiety as defined below or ω -A(C₁₋₆alkyl)CONH(C₁₋₆alkyl)-

5 wherein A is amino, mercapto, hydroxy or carboxyl;

R_2 is O (adenine 1-oxide derivatives), or is absent (adenine derivatives); or

10

R_1 and R_2 taken together form a 5-membered fused imidazole ring (1, N⁶-ethenoadenine derivatives), optionally substituted on the 4- or 5- positions of the etheno moiety with alkyl, aryl or aralkyl moieties as defined below;

15

R_3 is alkyl, aryl or aralkyl, alkylamino, arylamino or aralkylamino (NHR'); alkoxy, arylloxy or aralkyloxy (OR'); alkylthio, arylthio or aralkylthio (SR') as defined below; or ω -A(C₁₋₆alkyl)CONH(C₁₋₆alkyl)B- wherein A and B are independently amino, mercapto, hydroxy or carboxyl; or pharmaceutically acceptable esters, amides or salts thereof.


25

Thus the substituted derivatives of adenine include adenine 1-oxide; 1, N⁶-(4- or 5-substituted etheno) adenine; 6-substituted adenine; or 8-substituted aminoadenine, where R' of wherein the 6- or 8-HNR' groups are chosen from among: arylalkyl (C₁₋₆) groups with the aryl moiety optionally functionalized as described below; alkyl; and alkyl groups with functional groups therein, such as: 35 ([6-aminohexyl]carbamoylmethyl)-, and ω -acylated- amino(hydroxy, thiol and carboxy)alkyl(C₂₋₁₀)- and their ω -acylated-amino (hydroxy, thiol and carboxy) derivatives where the acyl group is chosen from among, but not limited to, acetyl, trifluoroacetyl, benzoyl, substituted-benzoyl, etc., or the carboxylic moiety is present as its ester or amide derivative, for example, the ethyl or methyl ester or its 45 methyl, ethyl or benzamido derivative. The ω -amino(hydroxy, thiol) moiety may be

alkylated with a C₁₋₄ alkyl group.

Likewise, B or B', or both, may be a pyrimidine with the general formula of Figure III, linked through the 1- position:

Formula III



wherein:

10 R₄ is hydrogen, hydroxy, mercapto, amino, cyano, aralkoxy, C₁₋₆ alkylthio, C₁₋₆ alkoxy, C₁₋₆ alkylamino or dialkylamino, the alkyl groups optionally linked to form a heterocycle;

 R₅ is hydrogen, acyl (e.g., acetoxy or benzoyl), C₁₋₆ alkyl, aroyl, optionally functionalized as defined below, C₁₋₅ alkanoyl, benzoyl, or sulphonate;

15 R₆ is hydroxy, mercapto, alkoxy, aralkoxy, C₁₋₆ alkylthio, amino, C₁₋₅ disubstituted amino, triazolyl, alkylamino or dialkylamino, where the alkyl groups are optionally linked to form a heterocycle or link to N³ to form an optionally substituted ring; or

 R₅ and R₆ taken together form a 5-membered fused imidazole ring between positions 3 and 4 of the pyrimidine ring (3,N⁴-ethenocytosine derivatives) optionally substituted on the 4- or 5- positions of the etheno moiety with alkyl, aryl or aralkyl moieties as defined below.

20 R₇ is hydrogen, hydroxy, cyano, nitro, alkenyl with the alkenyl moiety optionally linked through oxygen to form a ring optionally substituted on the carbon adjacent to the oxygen with alkyl or aryl groups, substituted alkynyl, halogen, alkyl, substituted alkyl,

perhalomethyl (e.g., CF_3), C_{2-6} alkyl, C_{2-3} alkenyl, or substituted ethenyl (e.g., allylamino, bromvinyl and ethyl propenoate, or propenoic acid), C_{2-3} alkynyl or substituted alkynyl; or together $\text{R}_6 - \text{R}_7$ may form a 5 or 6-membered saturated or unsaturated ring bonded through N or O at R_6 , such a ring may contain substituents that themselves contain functionalities; 5 provided that when R_8 is amino or substituted amino, R_7 is hydrogen; and

R_8 is hydrogen, amino or substituted amino, alkoxy, arylalkoxy, alkylthio, arylalkylthio, carboxamidomethyl, carboxymethyl, methoxy, methylthio, phenoxy or phenylthio; or pharmaceutically acceptable esters, amides or salts thereof.

In the general structures of Formula II and III above, the acyl groups 10 advantageously comprise alkanoyl or aroyl groups. The alkyl groups which may be straight or branched advantageously contain 1 to 8 carbon atoms, particularly 1 to 4 carbon atoms optionally substituted by one or more appropriate substituents, as described below. The aryl groups including the aryl moieties of such groups as aryloxy are preferably phenyl groups optionally substituted by one or more appropriate substituents, as described below. The 15 above-mentioned alkenyl and alkynyl groups advantageously contain 2 to 8 carbon atoms, particularly 2 to 6 carbon atoms, e.g., ethenyl or ethynyl, optionally substituted by one or more appropriate substituents as described below.

Appropriate substituents on the above-mentioned alkyl, alkenyl, alkynyl, and aryl groups are advantageously selected from halogen, hydroxy, C_{1-4} alkoxy, C_{1-4} alkyl, C_{6-10} 20 aryl, C_{7-12} arylalkyl, C_{7-12} arylalkoxy, carboxy, cyano, nitro, sulfonamido, sulfonate, phosphate, sulfonic acid, amino and substituted amino wherein the amino is singly or doubly substituted by a C_{1-4} alkyl, and when doubly substituted, the alkyl groups optionally being linked to form a heterocycle.

The compounds of the present invention encompass their pharmaceutically acceptable 25 esters, such as, but not limited to, acetyl and benzoyl esters. The esters may be made by reaction of the desired hydroxy compound with the appropriate acid, activated with carbonyldiimidazole, dicyclohexylcarbodiimide or other suitable condensing agent, or with an

acid anhydride or acid chloride with or without a basic catalyst such as a tertiary amine, quaternary ammonium salt or an inorganic base.

5 The compounds of the present invention also encompass their non-toxic pharmaceutically acceptable salts, such as, but not limited to, an alkali metal salt such as sodium or potassium; an alkaline earth metal salt such as manganese, magnesium or calcium; or an ammonium or tetraalkyl ammonium salt, i.e., NX_4^+ (wherein X is C_{1-4}).

Pharmaceutically acceptable salts are salts that retain the desired biological activity of the
10 parent compound and do not impart undesired toxicological effects. The present invention also encompasses the acylated prodrugs (e.g., esters) of the compounds disclosed herein. Those skilled in the art will recognize various synthetic methodologies which may be employed to prepare non-toxic pharmaceutically acceptable salts and acylated prodrugs of the compounds encompassed by Formulas I, IA and IB.

15 The compounds of the present invention are highly selective agonists of the $P2Y_2$ and/or $P2Y_4$ purinergic receptor; thus, they are useful in the treatment of mammals including humans suffering from chronic obstructive pulmonary diseases such as chronic bronchitis, PCD, cystic fibrosis, as well as prevention of pneumonia due to immobility. Furthermore, because of their general ability to clear retained mucus secretions and stimulate
20 ciliary beat frequency, the compounds of the present invention are also useful in the treatment of sinusitis, otitis media and nasolacrimal duct obstruction in mammals, including humans. Additionally, the compounds of the present invention are useful for treating mammals including humans with dry eye and retinal detachment.

 Though the compounds of the present invention are primarily concerned with
25 the treatment of human subjects, they may also be employed for the treatment of other mammalian subjects such as dogs and cats for veterinary purposes.

 The pharmaceutical utility of compounds of this invention are indicated by the inositol phosphate assay for $P2Y_2$ and other $P2Y$ receptor activity. This widely used assay,

as described in E. Lazarowski, et al., *Brit. J. Pharm.* 116, 1619-27 (1995), relies on the measurement of inositol phosphate formation as a measurement of activity of compounds activating receptors linked via G-proteins to phospholipase C.

The compounds of general Formulas I, IA, or IB may be administered orally, topically, parenterally, by inhalation or spray, intra-operatively, rectally, or vaginally in dosage unit formulations containing conventional non-toxic pharmaceutically acceptable carriers, adjuvants and vehicles. The term topically as used herein includes patches, gels, creams, ointments, or nose, ear or eye drops. The term parenteral as used herein includes subcutaneous injections, intravenous, intramuscular, intrasternal injection or infusion techniques. In addition, there is provided a pharmaceutical formulation comprising a compound of general Formulas I, IA or IB and a pharmaceutically acceptable carrier. One or more compounds of general Formulas I, IA or IB may be present in association with one or more non-toxic pharmaceutically acceptable carriers or diluents or adjuvants and, if desired, other active ingredients. One such carrier would be sugars, where the compounds may be intimately incorporated in the matrix through glassification or simply admixed with the carrier (e.g., lactose, sucrose, trehalose, mannitol) or other acceptable excipients for lung or airway delivery.

One or more compounds of general Formulas I, IA or IB may be administered separately or together, or separately or together with mucolytics such as DNase or acetylcysteine.

The pharmaceutical compositions containing compounds of general Formulas I, IA or IB may be in a form suitable for oral use, for example, as tablets, lozenges, aqueous or oily suspensions, dispersible powders or granules, emulsion, hard or soft capsules, or syrups or elixirs. Compositions intended for oral use may be prepared according to any method known to the art for the manufacture of pharmaceutical compositions and such compositions may contain one or more agents selected from the group consisting of sweetening agents, flavoring agents, coloring agents and preserving agents in order to

provide pharmaceutically elegant and palatable preparations. Tablets contain the active ingredient in admixture with non-toxic pharmaceutically acceptable excipients which are suitable for the manufacture of tablets. These excipients may be, for example, inert diluents, such as calcium carbonate, sodium carbonate, lactose, calcium phosphate or sodium phosphate; granulating and disintegrating agents, for example, corn starch, or alginic acid; binding agents, for example, starch, gelatin or acacia; and lubricating agents, for example magnesium stearate, stearic acid or talc. The tablets may be uncoated or they may be coated by known techniques to delay disintegration and absorption in the gastrointestinal tract and thereby provide a sustained action over a longer period. For example, a time delay material such as glyceryl monostearate or glyceryl distearate may be employed.

Formulations for oral use may also be presented as hard gelatin capsules wherein the active ingredient is mixed with an inert solid diluent, for example, calcium carbonate, calcium phosphate or kaolin, or as soft gelatin capsules wherein the active ingredient is mixed with water or an oil medium, for example, peanut oil, liquid paraffin or olive oil.

Aqueous suspensions contain the active materials in admixture with excipients suitable for the manufacture of aqueous suspensions. Such excipients are suspending agents, for example: sodium carboxymethylcellulose, methylcellulose and sodium alginate. Dispersing or wetting agents may be a naturally-occurring phosphatide or condensation products of an allylene oxide with fatty acids, or condensation products of ethylene oxide with long chain aliphatic alcohols, or condensation products of ethylene oxide with partial esters from fatty acids and a hexitol, or condensation products of ethylene oxide with partial esters derived from fatty acids and hexitol anhydrides. Those skilled in the art will recognize the many specific excipients and wetting agents encompassed by the general description above. The aqueous suspensions may also contain one or more preservatives, for example, ethyl, or n-propyl p-hydroxybenzoate, one or more coloring agents, one or more flavoring agents, and one or more sweetening agents, such as sucrose or saccharin.

Dispersible powders and granules suitable for preparation of an aqueous suspension by the addition of water provide the active ingredient in admixture with a dispersing or wetting agent, suspending agent and one or more preservatives. Suitable dispersing or wetting agents and suspending agents are exemplified by those already mentioned above. Additional excipients, for example, sweetening, flavoring, and coloring agents, may also be present.

Compounds of general Formulas I, IA or IB may be administered parenterally in a sterile medium. The drug, depending on the vehicle and concentration used, can either be suspended or dissolved in the vehicle. Advantageously, adjuvants such as local anaesthetics, preservatives and buffering agents can be dissolved in the vehicle. The sterile injectable preparation may be a sterile injectable solution or suspension in a non-toxic parentally acceptable diluent or solvent. Among the acceptable vehicles and solvents that may be employed are sterile water, saline solution, or Ringer's solution.

The compounds of general Formulas I, IA or IB may also be administered in the form of suppositories for ear, rectal or vaginal administration of the drug. These compositions can be prepared by mixing the drug with a suitable non-irritating excipient which is solid at ordinary temperatures but liquid at the body temperature and will therefore melt to release the drug. Such materials are cocoa butter and polyethylene glycols.

Solutions of compounds of general Formulas I, IA or IB may be administered by intra-operative installation.

Dosage levels of the order of from about 10^{-7} M to about 10^{-1} M, preferably in the range 10^{-5} to 10^{-1} M, are useful in the treatment of the above-indicated conditions. The amount of active ingredient that may be combined with the carrier materials to produce a single dosage form will vary depending upon the host treated and the particular mode of administration. It will be understood, however, that the specific dose level for any particular patient will depend upon a variety of factors including the activity of the specific compound employed, the age, body weight, general health, sex, diet, time of administration, route of

administration, and rate of excretion, drug combination and the severity of the particular disease undergoing therapy.

Compounds encompassed by the present invention may be prepared by condensation of a nucleoside mono-, di-, or triphosphate, activated with a condensing agent such as, but not limited to, carbonyldiimidazole or dicyclohexylcarbodiimide, with a second molecule of the same or a different mono-, di-, or triphosphate to form the desired dinucleotide polyphosphate; or a nucleoside phosphate, activated as above, may be condensed sequentially with a non-nucleoside mono-, di- or polyphosphate moiety, such as, but not limited to a monophosphate or pyrophosphate anion to yield the desired dinucleotide polyphosphate, the non-isolated intermediate in such a case being a mononucleotide polyphosphate; or a mono-, di- or polyphosphate moiety, activated as mentioned above, or in the form of an acid halide or other derivative reactive toward nucleophilic displacement, may be condensed sequentially with a nucleoside phosphate or polyphosphate to yield the desired dinucleotide polyphosphate; or the desired dinucleotide polyphosphate may be formed by modification a pre-formed dinucleotide polyphosphate by substitution or derivatization of a moiety or moieties on the purine, pyrimidine or carbohydrate ring. Nucleoside phosphates used as starting materials may be commercially available, or may be made from the corresponding nucleosides by methods well known to those skilled in the art. Likewise, where nucleosides are not commercially available, they may be made by modification of other readily available nucleosides, or by synthesis from heterocyclic and carbohydrate precursors by methods well known to those skilled in the art.

Those having skill in the art will recognize that the starting materials may be varied and additional steps employed to produce compounds encompassed by the present invention, as demonstrated by the following examples. In some cases protection of certain reactive functionalities may be necessary to achieve some of the above transformations. In general the need for such protecting groups will be apparent to those skilled in the art of organic synthesis as well as the conditions necessary to attach and remove such groups.

The invention is illustrated further by the following examples which are not to be construed as limiting the invention in scope or spirit to the specific procedures described in them.

5

Example 1

Preparation of P¹,P⁴-Di(uridine 5'-)P²,P³-methylenetetraphosphate

Methylenediphosphonic acid (Aldrich, 0.0088 g, 0.05 mmol) was dissolved in anhydrous DMF (0.5 mL) with the addition of tributylamine (24 μ L, 0.1 mmol). The solution was evaporated to dryness twice with anhydrous DMF (2 x 1 mL), the dried residue dissolved in anhydrous DMF (0.5 mL), and a solution of similarly-dried uridine 5'-monophosphomorpholidate 4-morpholine-N,N'-dicyclohexylcarboxamidinium salt (Sigma, 0.137 g, 0.2 mmol) in anhydrous DMF (0.5 mL) added. The reaction mixture was heated at 80-90° C for 7 h; then the solvent was removed by evaporation under reduced pressure. The residue was dissolved in water (2 mL) and applied to a column of DEAE cellulose (2.5 x 50 cm bed volume) in the bicarbonate form. The column was eluted with water, followed by a gradient of ammonium bicarbonate (0 - 0.33 M, in 900 mL). The progress of the elution was followed by monitoring absorbance of the eluate at 254 nm; the fraction eluting between 0.23 and 0.26 M was collected, evaporated to dryness and desalted by repeated evaporation with deionized water. The residue was dissolved in water (300 μ L) and purified in 50 μ L aliquots by semipreparative HPLC (Alltech PEI 5 μ , 10 x 250 mm, gradient 0 - 0.66-M ammonium bicarbonate, 5.0 mL/min, 20 min); peaks eluting at 8.7 - 9.0 min from each run were combined and lyophilized to yield the title compound (0.007 mmol, 14% yield, quantitated by comparison of its absorbance at λ_{max} 263 with that of a standard solution of uridine monophosphate). Chromatographic purity was 96.5% on an Alltech PEI column, gradient 0 - 0.66-M ammonium bicarbonate, 1.0 mL/min, 20 min., retention time 13.03 min. ¹H NMR in D₂O (δ ppm from tetramethylsilane): 2.39, (t, J = 21.5 Hz, 2H); 4.12 (m, 6H); 4.243 (m, 4H);

25

5.841 (d, $J = 7.9$ Hz, 2H); 5.847 (d, $J = 4.5$ Hz, 2H); 7.77, d, $J = 8.1$ Hz, 2H). ^{31}P NMR in D_2O (ppm from H_3PO_4) -10.2 to -10.7 (complex m, 2P); 7.8 to 8.4 (complex m, 2P).

5

Example 2

Preparation of P^1, P^4 -Di(uridine 5'- P^2, P^3 -difluoromethylenetetraphosphate)

10 The tributylammonium salt of difluoromethylenediphosphonic acid (as described in C. McKenna, et al., *J. Org. Chem.* 46, 4574-76 (1981) and D. Burton, et al., *J. Fluorine Chem.* 15, 263-66 (1980)) (0.014 g, 0.025 mmol), converted to the salt as described for methylenephosphonic acid was dissolved in a solution of uridine 5'-monophosphomorpholidate 4-morpholine- N, N' -dicyclohexylcarboxamidinium salt (Sigma, 15 0.034 g, 0.05 mmol) in anhydrous dimethyl sulfoxide (0.7 mL), and heated for 9 days at 50°C . The cooled reaction mixture was diluted with water and applied to a column of DEAE cellulose (2.5 x 50 cm bed volume) in the bicarbonate form. The column was eluted with water, followed by a gradient of ammonium bicarbonate (0 - 0.33 M, total volume 900 mL). The progress of the elution was followed by monitoring absorbance of the eluate at 254 nm. 20 The fraction eluting between 0.29 and 0.30 M was evaporated to dryness and desalted by repeated evaporations with deionized water to yield the title compound (0.0011 mmol, 4.4% yield, quantitated by comparison of its absorbance at λ_{max} 263 nm with that of a standard solution of uridine monophosphate). Chromatographic purity was 88.5% on an Alltech PEI column, gradient 0 - 0.66-M ammonium bicarbonate, 1.0 mL/min, 20 min., retention time 25 12.03 min. ^1H NMR in D_2O (δ ppm from tetramethylsilane): 4.05 - 4.085 (m, 6H); 4.18 - 4.20 (m, 4H); 5.80 (d, $J = 8.0$ Hz, 2H); 5.81 (d, $J = 4.5$ Hz, 2H); 7.77 (d, $J = 7.9$ Hz, 2H). ^{31}P NMR in D_2O (δ ppm from H_3PO_4) -10.63 (dd, $J = 18.3, 11.3$ Hz, 2P); - 5.83 (tdd, $J = 75, 18.3, 11.3$ Hz, 2P). ^{19}F NMR in D_2O : 73.406 (t, $J = 75.5$ Hz).

30

Example 3

Preparation of P¹,P⁴-Di(uridine 5'-P²,P³-imidotetraphosphate)

Tetrasodium imidodiphosphate (Sigma, 0.05 mmol) was dissolved in water 0.5 mL and applied to a column of Biorad AG-MP50 strong cation exchange resin (2 mL bed volume, 3 meq) in its tributylamine form. The column was eluted with water (~ 10 mL), the eluate lyophilized dried by evaporation with dry DMF. Treatment of uridine 5'-monophosphomorpholidate 4-morpholine-N,N'-dicyclohexylcarboxamidine salt (Sigma, 0.068 g, 0.1 mmol) with a solution of the tetrabutylammonium imidodiphosphate (0.05 mmol) in anhydrous DMF (1.0 mL) for 20 days at room temperature, and isolation essentially as described above yielded the title compound as the ammonium salt (1.6%). ¹H NMR in D₂O (δ ppm from tetramethylsilane): 4.07 - 4.09 (m, 6H); 4.17 - 4.22 (m, 4H); 5.79 (d, J = 8.1 Hz, 2H); 5.80 (d, J = 4.8 Hz, 2H); 7.78, d, J = 8.2 Hz, 2H). ³¹P NMR in D₂O (δ ppm from H₃PO₄) -10.82 (m, 4P); P-P coupling pattern similar to that of P¹,P⁴-di(adenosine 5'-tetraphosphate (Sigma) run under same conditions.

Example 4

Preparation of P¹,P⁴-Di(4-thiouridine 5'-tetraphosphate)

4-Thiouridine monophosphate sodium salt (Sigma, 25mg, 0.057 mmol) was dissolved in water 0.5 mL), applied to a column of Biorad AG-MP50 strong cation exchange resin (2 mL bed volume, 3 meq) in its tributylamine form, the column eluted with water (~ 10 mL) and the eluate lyophilized. The tributylammonium salt was dissolved in anhydrous DMF (0.5 mL) and carbonyldiimidazole (4.86 mg, 0.03 mmol) was added. The reaction mixture was set aside under nitrogen at room temperature for twelve days. The reaction mixture was evaporated to dryness under vacuum at room temperature, the residue dissolved in water (2 mL), and applied to a column of DEAE cellulose (2.5 x 50 cm bed volume) in the bicarbonate form. The column was eluted with water (~ 250 mL), then with a gradient of

ammonium bicarbonate (0 - 0.33 M, total volume 900 mL). This was followed by a gradient of 0.33 to 0.5 M ammonium bicarbonate over 400 mL. The progress of the elution was followed by monitoring absorbance of eluate at 280 nm. The fraction eluting between 0.336 and 0.339 M was evaporated to dryness and desalted by repeated evaporations with deionized water to yield the title compound (0.0045 mmol, 18% yield, quantitated by comparison of its absorbance at λ_{max} 332 nm with that of a standard solution of 4-thiouridine diphosphate). ^1H NMR in D_2O (δ ppm from tetramethylsilane): 4.09 - 4.11 (m, 6H); 4.18 - 4.24 (m, 4H); 5.76 (d, $J = 4.5$ Hz, 2H); 6.47 (d, $J = 7.7$ Hz, 2H); 7.67, d, $J = 8.2$ Hz, 2H). ^{31}P NMR in D_2O (δ ppm from H_3PO_4) -22.57 to -22.73 (m, 2P); -10.76 to -10.91 (m, 2P); P-P coupling pattern similar to that of P^1, P^4 -di(adenosine 5'-tetrphosphate (Sigma) run under same conditions.

Example 5

Preparation of P^1, P^5 -Di(uridine 5'-pentaphosphate)

A 100 ml round bottomed flask was charged with a DMF solution of uridine 5'-diphosphate tributylammonium salt (1.81 mmol, 10 ml) and carbonyldiimidazole (469 mg, 2.90 mmol) and the solution was stirred under N_2 for 2 hours. To this was added a DMF solution of uridine 5'-triphosphate tributylammonium salt (1.81 mmol, 10 ml) and the solution was stirred at 60°C for 24 hours. The solution was evaporated *in vacuo* and purified two times by column chromatography (DEAE Sephadex; $\text{H}_2\text{O} > 0.5\text{M NH}_4\text{HCO}_3$ gradient). The pure fractions were concentrated *in vacuo* at 35°C , and H_2O added and reevaporated ten times to obtain a white solid (200 mg). ^1H NMR in D_2O (δ ppm from tetramethylsilane): 4.0 (m, br, 6H), 4.1 (m, 4H), 5.7 (m, 4H), 7.7 (d, $J = 8.1$ Hz, 2H); ^{31}P NMR in D_2O (δ ppm from H_3PO_4) -22.3 (m, 3P), -10.6 (d, $J = 42.9$ Hz, 2P).

Example 6**Preparation of P¹,P⁴-Di(3,N⁴-ethenocytidine 5'-)tetraphosphate**

5 To a solution of P¹,P⁴-di(cytidine 5'-)tetraphosphate (reference 3, Table I; ammonium salt, 6 μ mol in 0.66 mL water) was added sodium bicarbonate (0.005 g, 60 μ mol) and the solution was lyophilized to remove ammonia. The residue was dissolved in a mixture of water and (0.20 mL) and chloroacetaldehyde solution (50% in water, 0.30 mL), and the reaction mixture set aside at room temperature for six days. The reaction mixture

10 was lyophilized, and the gummy residue partitioned between deuterium oxide (0.7 mL) and methylene chloride (1.5 mL). The ¹H NMR spectrum of the aqueous solution indicated that the ethenylation had progressed about 50%, while the ³¹P spectrum confirmed that the tetraphosphate chain remained intact. Additional chloroacetaldehyde solution (0.25 mL) was added to the NMR solution and the mixture set aside for a further ten days. The solution was

15 lyophilized, and the residue lyophilized again with deuterium oxide to remove exchangeable protons. The residue was partitioned between deuterium oxide and methylene chloride as before, and complete conversion to the ethenyl derivative confirmed by NMR spectroscopy. The deuterium oxide solution was applied to a column of DEAE cellulose (2.5 x 30 cm bed volume) in the bicarbonate form. The column was eluted with water (~250 mL), followed by

20 a gradient of 0 to 0.5 M ammonium bicarbonate over 1000 mL. The progress of the elution was followed by monitoring absorbance of the eluate at 280 nm. The fraction eluting between 0.29 and 0.32 M was evaporated to dryness and desalted by repeated evaporations with deionized water to yield the title compound (1.584 μ mol, 26.4% yield, quantitated by comparison of its absorbance at λ_{max} 273 nm with that of a standard solution of 3,N⁴-

25 ethenocytidine 5'-monophosphate). ¹H NMR in D₂O (δ ppm from tetramethylsilane): 4.123 (m, 6H); 4.258 (m, 4H); 5.986 (s, 2H); 6.92 (d, J = 8.1 Hz, 2H); 7.461 (s, 2H); 7.772 (s, 2H); 8.00 (d, J = 7.6 Hz, 2H). ³¹P NMR in D₂O (δ ppm from H₃PO₄) -22.474 (m, 2P); -10.650 (m,

2P); P-P coupling pattern closely similar to that of P¹,P⁴-di(adenosine 5'-tetraphosphate [Sigma]) run under same conditions.

5

Example 6(a)

P¹, P⁴-Di(imidazo[1,2-c]pyrimidin-5(6H)-one-2-(3-nitro)-phenyl-6-β-D-ribofuranoside 5'-)tetraphosphate, tetraammonium salt

P¹,P⁴-Di(cytidine 5'-)tetraphosphate, tetraammonium salt (100 mg, 0.117 mmol) (reference 3, Table I) was dissolved in water (10 mL) and flushed through a column of Dowex 50H⁺ resin (3g, pre-washed with methanol and water) and washed with 50 mL water. Tributylamine (1 mL) and dimethylformamide (DMF) (5 mL) were added and the solution was evaporated to an oil. The oil was dissolved in dry DMF (10 mL) and the evaporation cycle repeated twice. The final oil was dissolved in dry DMF (10 mL) and tributylamine (1.5 mL) to which was added α-bromo-3'-nitro-acetophenone (86 mg, 0.351 mmol). The reaction mixture was heated under nitrogen gas at 70°C for 20 hr, when more α-bromo-3'-nitro-acetophenone (50 mg, 0.205 mmol) was added. After heating for an additional 18 hr, the solvent was removed *in vacuo* and the residue purified by flash chromatography (DEAE Sephadex, 0>1.0M NH₄HCO₃) to obtain a yellow solid (13.5 mg): ¹H NMR (D₂O, TMS) δ 4.0-4.2 (m, br, 10H), 6.0 (d, J = 5.7 Hz, 2H), 6.4 (d, J = 8.1 Hz, 2H), 7.35 (t, J = 7.8 Hz, 2H), 7.5 (d, J = 8.1 Hz, 2H), 7.80 (m, 3H), 8.03 (s, 2H); ³¹P NMR (D₂O, H₃PO₄ std) δ -10.67 (m, 2P), -22.26 (m, 2P).

25

Example 7

P¹-(Thymidine-5'-)P⁴-(uridine-5'-)tetraphosphate (UP₂T)

A solution of uridine 5'-triphosphate (UTP) trisodium salt (ProBioSint, 5.86g, 0.01 mol) in water (5 mL) was passed through a column of BioRad AG-MP 50 strong cation

24

exchange resin in its tributylamine form (50 mL bed volume) and eluted with distilled water (about 300 mL). To this solution was added tributylamine (5 mL), and the suspension shaken until the pH of the aqueous fraction had risen to 8. The layers were separated and the aqueous solution evaporated to small volume, then lyophilized overnight. The residue was

5 dissolved in dry dimethylformamide (DMF, 20 mL) and the solvent evaporated at 0.1 mmHg. The dried tributylamine salt was made up to 100 mL with anhydrous acetone to yield a stock solution (0.1 M in UTP). Dicyclohexylcarbodiimide(DCC) (Baker, 0.1 g, 0.5 mmol) was added to an aliquot of the foregoing UTP solution (1.0 mL 0.1 mmol) and the solution stirred at room temperature for 30 min. The deposited dicyclohexylurea was removed by filtration,

10 the reaction mixture extracted with ether (10 mL), and the residue dissolved in dry deuterated dimethylsulfoxide (DMSO- d_6 , 0.3 mL). This solution of uridine 5'-cyclic metaphosphate (UcTP) was added to a solution of thymidine 5'-monophosphate (TMP, 0.064 g, 0.2 mmol) and tributylamine (0.2 mL) in DMSO- d_6 (0.3 mL) and set aside at 50° C for 24h. The reaction mixture was evaporated under high vacuum overnight, the residue dissolved in water

15 (1.0 mL), filtered to remove a little residual dicyclohexylurea, and separated by semipreparative ion-exchange chromatography (Hamilton PRP X-100 column, eluting with isocratic 1.0 M ammonium bicarbonate, 5 mL/min, 30 min, multiple injections of 100 μ L). The dinucleotide tetraphosphate eluted between 21 and 23 min; the product (11.1 % yield based on UTP) was quantitated by comparison of its ultraviolet absorption at λ_{max} 263 nm

20 with those of standards of UMP and TMP. 1H NMR D_2O , δ ppm from tetramethylsilane: 1.78 (s, 3H); 2.19-2.22 (m, 2H); 4.04-4.13 (m, 6H) 4.22-4.27 (m, 2H); 4.52 (m, partially obscured by D_2O); 4.74 (m, partially obscured by D_2O); 5.83 (d, J = 8.1 Hz, 1H); 5.84 (d, J = 5.0 Hz, 1H); 6.195 (t, J = 6.9 Hz, 1H); 7.61 (s, 1H); 7.82 (d, J = 8.1 Hz, 1H). ^{31}P NMR (D_2O , δ ppm from H_3PO_4) -22.71 (m, 2P); -10.97 (m, 2P).

Example 8**P¹-(Inosine 5'-)-P⁴-(uridine 5'-)tetraphosphate (UP₄I)**

Condensation of uridine 5'-cyclic trimetaphosphate (UcTP) and inosine 5'-monophosphate was carried out essentially as described above, except that the reaction mixture was stored at 25° C for five weeks prior to evaporation of the solvent. The residue was dissolved in water (1.0 mL), filtered, and separated in 150 µL aliquots by ion exchange chromatography on a Hamilton PRP X-100 column, eluting with isocratic 1.0 M ammonium bicarbonate, 5 mL/min, 30 min. The fractions eluting between 6 and 9 minutes were evaporated and lyophilized overnight to remove buffer. The dinucleotide tetraphosphate (8 % yield, 96% purity by HPLC (AUC)) was quantitated by comparison of its ultraviolet absorption at 260 nm with those of UMP and IMP at the same wavelength. ¹H NMR (D₂O, δ ppm from tetramethylsilane): 4.13 (m, 6H) 4.24-4.27 (m, 2H); 4.47 (m, partially obscured by D₂O); 5.78 (d, J = 7.9 Hz, 1H); 5.83 (d, J = 4.7 Hz, 1H); 6.003 (d, J = 5.7 Hz, 1H); 7.79 (d, J = 7.9 Hz, 1H); 8.10 (s, 1H); 8.37 (s, 1H). ³¹P NMR (D₂O, δ ppm from H₃PO₄) -22.43 (m, 2P); -10.72 (m, 2P).

Example 9**P¹-(4-Thiouridine 5'-)-P⁴-(uridine 5'-)tetraphosphate (UP₄(4-SH-U))**

4-Thiouridine monophosphate sodium salt (25mg, 0.057 mmol) was dissolved in water (0.5 mL), applied to a column of Biorad AG-MP50 strong cation exchange resin (2 mL bed volume, 3 meq) in its tributylamine form, the column eluted with water (~ 10 mL) and the eluate lyophilized. The resulting tributylamine salt of 4-thio-UMP was condensed with uridine 5'-cyclic trimetaphosphate (0.1 mmol) prepared by activation of UTP with dicyclohexylcarbodiimide (206 mg) essentially as described in Example 7 (72 h reaction time). After evaporation of the DMSO from the reaction mixture, the residue was dissolved in water (~1 mL) and separated in 200 µL aliquots by ion exchange chromatography on a

Hamilton PRP X-100 column, eluting with isocratic 1.0 M ammonium bicarbonate, 5 mL/min, monitoring the elution at 328 nm. The fractions eluting between 15 and 25 min were lyophilized to give the title compound (9.7% yield, 99.7 % pure by HPLC(AUC)), which was quantitated by comparison of its ultraviolet absorption at 332 nm with that of 4-thio-UMP at the same wavelength. ¹H NMR (D₂O, δ ppm from tetramethylsilane): 4.04 (m, partially overlapped by HOD); 4.14 (m, partially overlapped by HOD); 5.72 (m, 3H); 6.42 (d, J = 7.4 Hz); 7.55 (d, J = 7.6 Hz); 7.70 (d, J = 8.1 Hz). ³¹P NMR (D₂O, δ ppm from H₃PO₄) - 20.88 (m, 2P); -9.27 (m, 2P).

10 Examples 10-12 were prepared from uridine 5'-cyclic trimetaphosphate (0.1 mmol) and the relevant nucleoside 5'-monophosphate (0.2mmol) essentially as described in Example 7, except that hexane was used in place of ether to extract the excess of DCC from the reaction mixture.

15

Example 10

P¹-(Cytosine β-D-arabinofuranoside 5'-)P⁴-(uridine 5'-)tetraphosphate, (UP₄araC) (20mg); ¹H NMR (D₂O) δ 4.30-3.95 (m,10H), 5.99 (d,J=6.7Hz,1H), 6.08 (d,J=5.2Hz,1H), 7.82-7.77 (m,2H); ³¹P NMR (D₂O) δ -10.79 (m,2P), -22.52 (m,2P)

20

Example 11

P¹-(Uridine 5'-)-P⁴-(xanthosine 5'-)tetraphosphate (UP₄X) (27.7 mg); ¹H NMR (D₂O): δ 4.50-4.40 (m,10H), 5.80-5.70 (m,3H), 7.70 (d,J=8.0Hz,1H), 7.88 (s,1H,); ³¹P NMR (D₂O): δ -10.73 (m,2P), -22.41 (m,2P)

25

Example 12

P^1 -(2'-deoxyuridine 5'-)- P^4 -(uridine 5'-)tetraphosphate (UP_4dU) (40.6mg); 1H NMR (D_2O) δ 2.20-2.15 (m, 2H), 4.45-3.95 (m, 9H), 5.80-5.74 (m, 3H), 6.12 (t, $J=6.7$ Hz, 1H), 7.80-7.74 (m, 2H), ; ^{31}P NMR (D_2O) δ 2.9 (m, 2P), -8.9 (m, 2P)

5

Example 13

P^1 -(3'-Azido-3'-deoxythymidine 5'-)- P^4 -(uridine 5'-)tetraphosphate ($UP_4(AZT)$) and

Example 14, P^1, P^4 -Di(3'-azido-3'-deoxythymidine 5'-)tetraphosphate (AZT), P_4

10

3'-Azido-3'-deoxythymidine 5'-monophosphate (AZTMP) sodium salt (50 mg, 0.135 mmol) was dissolved in water (1 mL) and applied to a column of Biorad AG-MP50 strong cation exchange resin in its tributylamine form. The column was eluted with water (~ 10 mL) and the eluate lyophilized. The resulting tributylamine salt was condensed with uridine

15 5'-cyclic trimetaphosphate (0.1 mmol) prepared by activation of UTP with dicyclohexylcarbodiimide (206 mg) essentially as described in Example 7. The residue after evaporation of the reaction mixture was dissolved in water (1.0 mL), passed through a 0.45 μ syringe filter to remove a little solid, and the filtrate subjected to preparative HPLC on a Hamilton PRP X-100 column, eluting with an isocratic mixture of 1.0 M ammonium

20 bicarbonate (75%) and methanol (25%), (4 mL/min). The fraction eluting between 5 and 8 min was lyophilized to yield $UP_4(AZT)$ (7.9%), quantitated by comparison of its ultraviolet absorption at 264 nm with those of UMP and TMP at the same wavelength. 1H NMR D_2O , δ ppm from tetramethylsilane: 1.78 (s, 3H); 2.30-2.34 (m, 2H); 4.07-4.14 (m, 6H) 4.22-4.29 (m, 2H); 4.52 (m, 1H); 5.82 (d, $J = 4.4$ Hz, 1H); 5.84 (d, $J = 8.1$ Hz, 1H); 6.12 (t, $J = 7$ Hz, 1H); 7.62 (s, 1H); 7.81 (d, $J = 8.1$ Hz, 1H). ^{31}P NMR (D_2O , δ ppm from H_3PO_4) -22.51 (m, 2P); -11.06 (m, 1P); -10.81 (m, 1P).

25

Collection and lyophilization of the fractions eluting between 25 to 40 minutes from the same reaction mixture yielded P^1, P^4 -Di(3'-azido-3'-deoxythymidine 5'-)tetraphosphate (3%), quantitated by comparison of its ultraviolet spectrum at 266 nm with that of TMP. 1H NMR D_2O , δ ppm from tetramethylsilane: 1.80 (s, 6H); 2.31-2.35 (m, 4H); 4.09-4.11 (m, 6H)
 5 4.46-4.47 (m, 2H); 6.12 (t, $J = 7$ Hz, 2H); 7.63 (s, 2H). ^{31}P NMR (D_2O , δ ppm from H_3PO_4) - 22.47 (m, 2P); -11.35 (m, 2P).

Example 14

P^1, P^6 -Di(uridine-5'-)hexaphosphate (U_2P_6)

10

The dinucleotide hexaphosphate (6.97%) was formed by reaction of uridine cyclic trimetaphosphate with uridine 5'-triphosphate under similar conditions. 1H NMR D_2O , δ ppm from tetramethylsilane: 4.06-4.19 (m, 6H); 4.21-4.4.26 (m, 4H); 5.78 (d, $J = 8.2$ Hz, 2H); 5.81 (d, $J = 5.4$ Hz, 2H); 7.78 (d, $J = 8.1$ Hz, 1H), ^{31}P NMR (D_2O , δ ppm from H_3PO_4) -
 15 22.41 (m, 4P); -10.89 (m, 2P).

Examples 15 and 16

2'(3')-Benzoyl- P^1, P^4 -di(uridine 5'-)tetraphosphate (Example 15)

and P^1, P^4 -Di(2'(3')-benzoyl uridine 5'-)tetraphosphate (Example 16)

20 Benzoic acid (61.7 mg, 0.505 mmol) and 1,1-carbonyldiimidazole (81.8 mg, 0.505 mmol) were combined in anhydrous DMF (1 mL) and stirred at ambient temperature for 1 hour. P^1, P^4 -Di(uridine 5'-)tetraphosphate (97 mg, 0.102 mMol) in anhydrous DMF (2 mL) was added and the mixture stirred at ambient temperature for 4 hours. The temperature was increased to 35°C and stirring continued for 6 days. The reaction mixture was evaporated to
 25 dryness, dissolved in water, applied to a Sephadex DEAE A25 column (2.5 x 20 cm) and eluted with a ammonium bicarbonate gradient (0 to 0.3M, 400 mL total volume) followed by

isocratic ammonium bicarbonate (0.5M, 500 mL). Two fractions were collected, evaporated to dryness, then repeatedly co-evaporated with water to remove ammonium salts. The material eluting earlier was identified as the monobenzoyl ester: ^1H NMR (D_2O) δ 4.0-4.23 (m,9H), 5.35-5.45 (m,1H), 5.65-5.85 (m,3H), 5.98-6.02 (m,1H), 7.32-7.95 (m,7H); ^{31}P NMR (D_2O) δ -10.70 (m,2P), -22.28 (m,2P): the material eluting later was identified as the dibenzoyl ester: ^1H NMR (D_2O) δ 4.05-4.40 (m,8H), 5.30-6.05 (m,6H), 7.2-7.95 (m,12H); ^{31}P NMR (D_2O) δ -10.70 (m,2P), -22.45 (m,2P)

Examples 17, 18, 19 and 20

$\text{P}^1\text{-(2'-deoxyguanosine 5')P}^4\text{-(uridine 5')tetrphosphate (UP}_4\text{dG)}$ (Example 17);
 $\text{P}^1\text{-(2'-deoxyadenosine 5')P}^4\text{-(uridine 5')tetrphosphate (UP}_4\text{dA)}$ (Example 18);
 $\text{P}^1\text{-(2'-deoxyinosine 5')P}^4\text{-(uridine 5')tetrphosphate (UP}_4\text{dI)}$ (Example 19) and
 $\text{P}^1\text{-(2'-deoxycytidine 5')P}^4\text{-(uridine 5')tetrphosphate (UP}_4\text{dC)}$ (Example 20)

Uridine 5'-triphosphate (5.0 g) in water (18 mL) was passed through a column of Dowex 50 H^+ , and tributylamine (3.0 g) added to the eluent. The mixture was concentrated to an oil, dried by evaporation with dry DMF and redissolved in dry DMF (18 mL). Dicyclohexylcarbodiimide (DCC, 3.5 g) was added, the solution stirred at room temperature for 30 min, and the precipitate removed by filtration. Hexane (70 mL) was added to the filtrate, the bottom layer separated and washed again with hexane (70 mL) to complete the removal of DCC. This solution of uridine 5'-cyclic metaphosphate (UcTP) was used in the following experiments:

Example 17**P¹-(2'-deoxyguanosine 5'-)P⁴-(uridine 5'-)tetraphosphate (UP₄dG)**

2'-Deoxyguanosine 5'-monophosphate (d-GMP, Sigma, 500 mg) was dissolved in
5 DMF (4.5mL), tributylamine (1.0 mL) added, and the solution concentrated to an oil under
vacuum. One third of the solution of uridine 5'-cyclic metaphosphate (UcTP, above) was
added and the solution heated at 40 °C for 24 h. The solution was evaporated to an oil,
dissolved in water (10 mL) and applied to a column of Sephadex DEAE (350 mL in a 4.5 x
22 cm column) in its bicarbonate form, pre-equilibrated with 0.25 M ammonium bicarbonate.
10 The column was eluted successively with 0.25, 0.30, 0.35, 0.40, and 0.50 M ammonium
bicarbonate. The elution was monitored by HPLC (SynchroPak AX-300, 75% 0.50 M
KH₂PO₄, 25% MeCN 1.0 mL/min, UV 254 nm), and the fraction containing UP₄dG was
concentrated to a solid, then co-evaporated 6-7 times with water to yield the ammoniu salt of
the dinucleotide as an orange-yellow solid (140 mg, estimated purity by HPLC (AUC) 94%).

15

Example 18**P¹-(2'-deoxyadenosine 5'-)P⁴-(uridine 5'-)tetraphosphate (UP₄dA)**

Reaction of 2'-deoxyadenosine 5'-monophosphate (d-AMP, Sigma, 500 mg) with
20 uridine 5'-cyclic metaphosphate essentially as described above gave UP₄dA as the solid white
ammonium salt (140 mg, HPLC purity as above, 99%).

Example 19**P¹-(2'-deoxyinosine 5'-)P⁴-(uridine 5'-)tetraphosphate (UP₄dI)**

25

2'-Deoxyinosine 5'-monophosphate (d-IMP, Sigma, sodium salt 1.0 g) is converted to
the free acid form with Dowex 50 (H⁺) resin as described for UTP above. The eluent is

neutralized with tributylamine (2.0 mL) and the mixture concentrated to an oil under vacuum. The resulting tributylamine salt is dried by evaporation with DMF, the residue dissolved in DMF (4.5 mL) and treated with uridine 5'-cyclic trimetaphosphate essentially as described above to yield P¹-(2'-deoxyinosine 5'-)-P⁴-(uridine 5'-)tetraphosphate.

5

Example 20

P¹-(2'-deoxycytidine 5'-)-P⁴-(uridine 5'-)tetraphosphate (UP₄dC)

2'-Deoxycytidine 5'-monophosphate (d-CMP, Sigma, 500 mg) treated essentially as above yielded P¹-(2'-deoxycytidine 5'-)-P⁴-(uridine 5'-)tetraphosphate as a white solid (130

10 mg) estimated HPLC purity 82%.

Example 21

Pharmacological activity as measured by the inositol phosphate assay

15

The compounds of Examples 1 - 20 were tested for their ability to elicit P2Y₁, P2Y₂, P2Y₄ and P2Y₆ receptor activity using the inositol phosphate assay as described by E. Lazarowski, et al., *Brit. J. Pharm.* 116, 1619-27 (1995). The results are summarized in Table II below.

20

25

TABLE II
DINUCLEOTIDE ACTIVITY SUMMARY

5

EC₅₀'s (μmol)

Example	P2Y1	P2Y2	P2Y4	P2Y6
1	IA	WEAK	11.1 (60%)	IA
2	IA	5.71	1.0 (80%)	WEAK
3	3.67	0.63	1.19	2.56
4	IA	0.02	n/a	0.05 (20%)
5	31.2 (55%)	3.8 (80%)	2.87	92.64
6	WEAK	0.46	19.8 (75%)	IA
6a	nd	0.3	0.06	0.87
7	IA	0.11	0.2	0.88
8	IA	0.11	0.15	0.8
9	>100	0.37	4.16	1.79
10	IA	0.19	0.32(65%)	1.3
11	IA	0.13	0.38(75%)	3.39
12	IA	0.1	0.39	0.92
13	nd	0.2	0.13	0.54
14	nd	10.7	14.7	3.5
15	IA	0.64	3.72	9.71
16	IA	0.79	0.73	5.33

IA Response < 2-fold basal

WEAK EC₅₀ > 100μmol

10 (XX%) Percentage response of same study positive control


nd not determined

The invention, and the manner and process of making and using it, are now described in such full, clear, concise and exact terms as to enable any person skilled in the art to which it pertains, to make and use the same. It is to be understood that the foregoing describes
5 preferred embodiments of the present invention and that modifications may be made therein without departing from the spirit or scope of the present invention as set forth in the claims. To particularly point out and distinctly claim the subject matter regarded as invention, the following claims conclude this specification.

WHAT IS CLAIMED IS:

1. A compound of Formula I:

5

Formula I

wherein:

10

X is oxygen, methylene, difluoromethylene, imido;

n = 0, 1 or 2;

m = 0, 1 or 2;

15

n + m = 0, 1, 2, 3 or 4; and

B and B' are each independently a purine residue or a pyrimidine residue linked through the 9- or 1- position, respectively;

20

Z = OH or N₃;

Z' = OH or N₃;

Y = H or OH;

Y' = H or OH;

provided that when Z is N₃, Y is H or when Z' is N₃, Y' is H; and further provided that the

25

compounds of Table I are excluded;

TABLE I
DINUCLEOTIDES

5

Np ₂ N	Np ₂ N'	Np ₃ N	Np ₃ N'	Np ₄ N	Np ₄ N'
Ap ₂ A	Ap ₂ NAD	Up ₃ U	Ap ₃ T	Up ₄ U	Ap ₄ U
Gp ₂ G	Ap ₂ TAD	Ap ₃ A	m ⁷ Gp ₃ G	Ap ₄ A	Ap ₄ C
m ⁷ Gp ₂ m ⁷ G	Ap ₂ C-NAD	Xp ₃ X	m ^{2,2'} Gp ₃ G	Cp ₄ C	Ap ₄ G
	Ap ₂ C-PAD	m ⁷ Gp ₃ m ⁷ G	m ^{2,7} Gp ₃ G	Gp ₄ G	Gp ₄ U
	Ap ₂ BAD	Gp ₃ G		Xp ₄ X	Gp ₄ C
	m ⁷ Gp ₂ G			Dp ₄ D	Up ₄ C
	Up ₂ U			eAp ₄ eA	Ap ₄ T
				m ⁷ Gp ₄ m ⁷ G	m ⁷ Gp ₄ G
					m ^{2,7} Gp ₄ G
					m ^{2,2,7} Gp ₄ G

Np ₂ N	Np ₂ N'	Np ₃ N	Np ₃ N'	Np ₄ N
Ap ₂ A	Ap ₂ T	Ap ₃ A	Ap ₃ T	Ap ₄ A

AppZppA	DppZppD	ApZppZpA	ApSpZpSpA
Z	Z	Z	Z
CH ₂	CH ₂	CH ₂	CHF
CH ₂ CH ₂	CH ₂ CH ₂	CH ₂ CH ₂	CF ₂
CHF	CHF	CHF	O
CF ₂	CF ₂	CF ₂	
CHCl	CHCl	CHCl	
CCl ₂	CCl ₂	CCl ₂	

10


- A = Adenosine
 U = Uridine
 G = Guanosine
 T = Thymidine
 X = Xanthosine
 TAD = Tiazofurin
 BAD = Benzamide riboside
 D = 2,6-Diaminopurine
- eA = Ethenoadenosine
 m⁷G = 7-Methylguanosine
 m^{2,7}G = 2,7-Dimethylguanosine
 m^{2,2,7}G = 2,2,7-Trimethylguanosine
 NAD = nicotinamide riboside
 C-NAD = C-nicotinamide riboside
 C-PAD = C-picolinamide riboside
 N = Nucleoside

20

or pharmaceutically acceptable esters or salts thereof.

2. A compound according to claim 1 of Formula IA:

Formula IA



wherein:

- 10 $X=O$;
 $n+m=1$ or 2 ;
 Z, Z', Y and $Y'=OH$;
 B and B' are uracil, thymine, cytosine, guanine, adenine, xanthine, hypoxanthine or
as defined in Formulas II and III; or

- 15 $X=O$;
 $n+m=3$ or 4 ;
 Z, Z', Y and $Y'=OH$;
 $B=uracil$;
20 B' is uracil, thymine, cytosine, guanine, adenine, xanthine, hypoxanthine or
as defined in Formulas II and III; or

- $X=O$;
 $n+m=1$ or 2 ;
25 Z, Y and $Y'=OH$;
 $Z'=H$;
 $B=uracil$;
 B' is uracil, thymine, cytosine, guanine, adenine, xanthine, hypoxanthine or
as defined in Formulas II and III; or

- 30 $X=O$;
 $n+m=0, 1$ or 2 ;
 Z and $Y=OH$;
 $Z'=N_3$;
35 $Y'=H$;
 $B=uracil$;
 $B'=thymine$; or

$X=O$;

$n+m=0, 1 \text{ or } 2$;
 $Z \text{ and } Z'=N_3$;
 $Y \text{ and } Y'=H$;
 $B \text{ and } B'=\text{thymine}$; or


$X=CH_2, CF_2 \text{ or } NH$;

$n \text{ and } m=1$;

$Z, Z', Y \text{ and } Y'=OH$;

$B \text{ and } B'$ are uracil, thymine, cytosine, guanine, adenine, xanthine, hypoxanthine or as defined in Formulas II and III;

Formula II



wherein

R_1 is C_{1-8} alkyl; phenyl or phenyloxy (which may be substituted with halogen; hydroxy; C_{1-4} alkoxy; C_{1-4} alkyl; C_{6-10} aryl; carboxy; cyano; nitro; sulfonamido; sulfonate; phosphate; sulfonic acid; amino or substituted amino, wherein the amino is singly or doubly substituted by a C_{1-4} alkyl and when doubly substituted, the alkyl groups may be linked to form a heterocycle); or or $\omega\text{-A}(C_{1-6}\text{alkyl})\text{CONH}(C_{1-6}\text{alkyl})\text{-}$ wherein A is amino, mercapto, hydroxy or carboxyl; or

R_2 is O or is absent; or

R_1 and R_2 taken together may form a 5-membered fused imidazole ring (which may

be substituted on the 4- or 5- positions of the etheno moiety with C₁₋₄alkyl; phenyl or phenyloxy, which may be substituted with halogen; hydroxy; C₁₋₄alkoxy; C₁₋₄alkyl; C₆₋₁₀aryl; carboxy; cyano; nitro; sulfonamido; sulfonate; or phosphate;

5 ₁₀aryl; carboxy; cyano; nitro; sulfonamido; sulfonate; or phosphate;

sulfonic acid; amino or substituted amino, wherein the amino is singly or doubly substituted by a C₁₋₄ alkyl and when doubly substituted, the alkyl groups may be linked to form a

10 heterocycle or C₇₋₁₂arylalkyl);

R₃ is C₁₋₈alkyl; phenyl or phenyloxy (which may be substituted with halogen;

15 hydroxy; C₁₋₄alkoxy; C₁₋₄alkyl; C₆₋₁₀aryl; carboxy; cyano; nitro; sulfonamido; sulfonate;

phosphate; sulfonic acid; amino or substituted amino, wherein the amino is singly or doubly

substituted by a C₁₋₄ alkyl and when doubly substituted, the alkyl groups may be linked to


20 form a heterocycle); or C₇₋₁₂arylalkyl; C₁₋₄alkylamino, phenylamino or C₇₋₁₂arylalkylamino,

C₁₋₄alkoxy, or C₇₋₁₂arylalkyloxy; C₁₋₄alkylthio, phenylthio or C₇₋₁₂arylalkylthio; or

ω-A(C₁₋₆alkyl)CONH(C₁₋₆alkyl)B- wherein A and B are independently amino, mercapto, hydroxy or carboxyl;

25

Formula III



wherein:

R_4 is hydrogen, hydroxy, mercapto, amino, cyano, C_{7-12} arylalkoxy, C_{1-6} alkylthio, C_{1-6} alkoxy, C_{1-6} alkylamino or diC_{1-4} alkylamino, wherein the alkyl groups may be linked to form a heterocycle;

R_5 is hydrogen, acetoxy or benzoyl, C_{1-6} alkyl, phenoxy, C_{1-5} alkanoyl, benzoyl, or sulphonate;

R_6 is hydroxy, mercapto, C_{1-4} alkoxy, C_{7-12} arylalkoxy, C_{1-6} alkylthio, amino, C_{1-5} disubstituted amino, triazolyl, C_{1-6} alkylamino or diC_{1-4} alkylamino, where the alkyl groups may be linked to form a heterocycle or link to N^3 to form a substituted ring; or


R_5 and R_6 taken together form a 5-membered fused imidazole ring between positions 3 and 4 of the pyrimidine ring ($3,N^4$ -ethenocytosine derivatives) which may be substituted on the 4- or 5- positions of the etheno moiety with C_{1-4} alkyl; phenyl or phenoxy, which may be substituted with halogen; hydroxy; C_{1-4} alkoxy, C_{1-4} alkyl; C_{6-10} aryl; carboxy; cyano; nitro; sulfonamido; sulfonate or phosphate; sulfonic acid; amino or substituted amino, wherein the amino is singly or doubly substituted by a C_{1-4} alkyl and when doubly substituted, the alkyl groups may be linked to form a heterocycle; or C_{7-12} arylalkyl;

R_7 is hydrogen, hydroxy, cyano, nitro, or C_{2-8} alkenyl; wherein the alkenyl moiety may be linked through oxygen to form a ring which may be substituted on the carbon adjacent to the oxygen with C_{1-4} alkyl, phenyl, substituted C_{2-8} alkynyl, halogen, C_{1-4} alkyl, substituted C_{1-4} alkyl, CF_3 , C_{2-6} alkyl, C_{2-3} alkenyl, allylamino, bromvinyl, ethyl propenoate, propenoic acid, C_{2-3} alkynyl, substituted C_{2-3} alkynyl; or together $R_6 - R_7$ may form a 5 or 6-membered saturated or unsaturated ring bonded through N or O at R_6 , such ring may contain substituents that themselves contain functionalities; provided that when R_6 is amino or substituted amino, R_7 is hydrogen;

R_8 is hydrogen, amino or diC_{1-4} alkylamino, C_{1-4} alkoxy, C_{7-12} arylalkoxy, C_{1-4} alkylthio, C_{7-12} arylalkylthio, carboxamidomethyl, carboxymethyl, methoxy, methylthio, phenoxy or phenylthio.

3. A compound according to claim 1 of Formula IB:

Formula IB



5 wherein:

X is oxygen, methylene, difluoromethylene or imido;


n = 0 or 1;

m = 0 or 1;

n + m = 0, 1 or 2; and

10 B and B' are each independently a purine residue, as in Formula II, or a pyrimidine residue, as in Formula III, linked through the 9- or 1- position, respectively;

Formula II



15 wherein

R₁ is C₁₋₈alkyl; phenyl or phenoxy (which may be substituted with halogen;

hydroxy; C₁₋₄alkoxy; C₁₋₄alkyl; C₆₋₁₀aryl; carboxy; cyano; nitro; sulfonamido; sulfonate;

20

phosphate; sulfonic acid; amino or substituted amino, wherein the amino is singly or doubly


substituted by a C₁₋₄ alkyl and when doubly substituted, the alkyl groups may be linked to
form a heterocycle); or or ω -A(C₁₋₆alkyl)CONH(C₁₋₆alkyl)- wherein A is amino, mercapto,
5 hydroxy or carboxyl; or

R₂ is O or is absent; or

R₁ and R₂ taken together may form a 5-membered fused imidazole ring (which may
be substituted on the 4- or 5- positions of the etheno moiety with C₁₋₄alkyl; phenyl or
15 phenoxy, which may be substituted with halogen; hydroxy; C₁₋₄alkoxy; C₁₋₄alkyl;
C₆₋₁₀ aryl; carboxy; cyano; nitro; sulfonamido; sulfonate; or phosphate;

sulfonic acid; amino or substituted amino, wherein the amino is singly or doubly substituted
20 by a C₁₋₄ alkyl and when doubly substituted, the alkyl groups may be linked to form a
heterocycle); or C₇₋₁₂arylalkyl;

R₃ is C₁₋₈alkyl; phenyl or phenoxy (which may be substituted with halogen;
hydroxy; C₁₋₄alkoxy; C₁₋₄alkyl; C₆₋₁₀aryl; carboxy; cyano; nitro; sulfonamido; sulfonate;
phosphate; sulfonic acid; amino or substituted amino, wherein the amino is singly or doubly
30 substituted by a C₁₋₄ alkyl and when doubly substituted, the alkyl groups may be linked to
form a heterocycle); or C₇₋₁₂arylalkyl; C₁₋₄alkylamino, phenylamino or
C₇₋₁₂ arylalkylamino, C₁₋₄alkoxy, or C₇₋₁₂arylalkoxy; C₁₋₄alkylthio, phenylthio or
C₇₋₁₂ arylalkylthio; or ω -A(C₁₋₆alkyl)CONH(C₁₋₆alkyl)B- wherein A and B are independently
35 amino, mercapto, hydroxy or carboxyl;

Formula III

wherein:

- 5 R_4 is hydrogen, hydroxy, mercapto, amino, cyano, C_{7-12} arylalkoxy, C_{1-6} alkylthio, C_{1-6} alkoxy, C_{1-6} alkylamino or diC_{1-4} alkylamino, wherein the alkyl groups may be linked to form a heterocycle;

R_5 is hydrogen, acetoxy or benzoyl, C_{1-6} alkyl, phenyloxy, C_{1-5} alkanoyl, benzoyl, or sulphonate;

- 10 R_6 is hydroxy, mercapto, C_{1-4} alkoxy, C_{7-12} arylalkoxy, C_{1-6} alkylthio, amino, C_{1-5} disubstituted amino, triazolyl, C_{1-6} alkylamino or diC_{1-4} alkylamino, where the alkyl groups may be linked to form a heterocycle or link to N^3 to form a substituted ring;

- R_5 and R_6 taken together form a 5-membered fused imidazole ring between positions 3 and 4 of the pyrimidine ring (3, N^4 -ethenocytosine derivatives) which may be substituted on the 4- or 5- positions of the etheno moiety with C_{1-4} alkyl; phenyl; phenyloxy or 15 C_{7-12} arylalkyl, which may be substituted with halogen; hydroxy; C_{1-4} alkoxy; C_{1-4} alkyl; C_6 , $_{10}$ aryl; carboxy; cyano; nitro; sulfonamido; sulfonate; phosphate; sulfonic acid; amino or substituted amino, wherein the amino is singly or doubly substituted by a C_{1-4} alkyl and when doubly substituted, the alkyl groups may be linked to form a heterocycle;

- 20 R_7 is hydrogen, hydroxy, cyano, nitro, or C_{2-8} alkenyl; wherein the alkenyl moiety may be linked through oxygen to form a ring which may be substituted on the carbon adjacent to the oxygen with C_{1-4} alkyl, phenyl, substituted C_2 .

alkynyl, halogen, C₁₋₄alkyl, substituted C₁₋₄alkyl, CF₃, C₂₋₆ alkyl, C₂₋₃ alkenyl, allylamino, bromvinyl, ethyl propenoate, propenoic acid, C₂₋₃ alkynyl, substituted C₂₋₃alkynyl; or together R₆ - R₇ may form a 5 or 6-membered saturated or unsaturated ring bonded through N or O at R₆, such ring may contain substituents that themselves contain functionalities; provided that when R₈ is amino or substituted amino, R₇ is hydrogen;

R₈ is hydrogen, amino or di-C₁₋₄amino, C₁₋₄alkoxy, C₇₋₁₂arylalkoxy, C₁₋₄alkylthio, C₇₋₁₂arylalkylthio, carboxamidomethyl, carboxymethyl, methoxy, methylthio, phenoxy or phenylthio.

4. A compound according to claims 2 or 3 where the acyl groups of Formulas II and III comprise alkyl or aryl groups, the alkyl groups having 1 to 4 carbon atoms, and the aryl groups including the aryl moieties of such groups as aryloxy are phenyl groups, where the alkyl and aryl groups are substituted with substituents selected from the group consisting of halogen, hydroxy, C₁₋₄ alkoxy, C₁₋₄ alkyl, C₇₋₁₂ aryl, C₇₋₁₂ arylalkoxy, carboxy, cyano, nitro, sulfonamido, sulfonate, phosphate, sulfonic acid, amino and substituted amino wherein the amino is singly or doubly substituted by a C₁₋₄ alkyl, and when doubly substituted, the alkyl groups are linked to form a heterocycle.

5. A compound according to claims 1-3, wherein B and B' are uracil attached at the N-1 position to the ribosyl moiety, and wherein the total of m + n equals 3 or 4 when X is oxygen.

6. A compound according to claims 1-3, wherein the ribosyl moieties are in the D- configuration.

7. A compound according to claim 1-3, wherein the ribosyl moieties are in the L- configuration.

8. A compound according to claim 1-3, wherein the ribosyl moieties are in the D- and L- configuration.

- 5 9. P^1, P^4 -Di(uridine 5'-) P^2, P^3 -methylenetetrphosphate.
10. P^1, P^4 -Di(uridine 5'-) P^2, P^3 -difluoromethylenetetrphosphate.
11. P^1, P^4 -Di(uridine 5'-) P^2, P^3 -imidotetrphosphate.
- 10 12. P^1, P^4 -Di(4-thiouridine 5'-)tetrphosphate.
13. P^1, P^5 -Di(uridine 5'-)pentaphosphate.
- 15 14. P^1, P^4 -Di(3,N⁴-ethenocytidine 5'-)tetrphosphate.

15. A pharmaceutical composition comprising a compound of Formulas I, IA or IB as described in claims 1-3 or a pharmaceutically acceptable salt thereof together with a pharmaceutically acceptable carrier therefor.

20

16. A method of treating chronic obstructive pulmonary diseases in a mammal by administering an effective chronic obstructive pulmonary disease treatment amount of a compound of Formulas I, IA or IB as described in claims 1-3.

25

17. A method of treating sinusitis, otitis media or nasolacrimal duct obstruction in a mammal by administering an effective mucus secretion clearing amount of a compound of Formulas I, IA or IB as described in claims 1-3.

18. A method of treating dry eye in a mammal by administering an effective dry eye treatment amount of a compound of Formula I, IA or IB as described in claims 1-3.

5 19. A method of treating retinal detachment in a mammal by administering an effective retinal detachment treatment amount of a compound of Formula I as described in claim 1.

20. A method of facilitating sputum induction in a mammal by administering an amount of a compound of Formula I as described in claim 1 effective to facilitate sputum induction.

21. A method of facilitating expectoration in a mammal by administering an amount of a compound of Formula I as described in claim 1 effective to facilitating expectoration.

22. A compound selected from the group consisting of:

P^1 -(Thymidine 5'-)- P^4 -(uridine 5'-) tetraphosphate (UP_4T),
 P^1 -(Inosine 5'-)- P^4 -(uridine 5'-)tetraphosphate (UP_4I),
20 P^1 -(4-Thiouridine 5'-)- P^4 -(uridine 5'-)tetraphosphate ($UP_4(4-SH-U)$),
 P^1 -(Cytosine β -D-arabinofuranoside 5'-)- P^4 -(uridine 5'-)tetraphosphate (UP_4araC),
 P^1 -(Uridine 5'-)- P^4 -(xanthosine 5'-)tetraphosphate (UP_4X),
 P^1 -(2'-deoxyuridine 5'-)- P^4 -(uridine 5'-)tetraphosphate (UP_4dU),
25 P^1 -(3'-Azido-3'-deoxythymidine 5'-)- P^4 -(uridine 5'-)tetraphosphate ($UP_4(AZT)$),
 P^1, P^4 -Di(3'-azido-3'-deoxythymidine 5'-)tetraphosphate ($((AZT)_2P_4)$),

5

P^1, P^6 -Di(uridine 5'-)hexaphosphate (U_2P_6),
2'(3')-Benzoyl- P^1, P^4 -di(uridine 5'-)tetraphosphate,
 P^1, P^4 -Di(2'(3')-benzoyluridine 5'-)tetraphosphate,
 P^1 -(2'-deoxyguanosine 5'-)- P^4 -(uridine 5'-)tetraphosphate (UP_4dG),
 P^1 -(2'-deoxyadenosine 5'-)- P^4 -(uridine 5'-)tetraphosphate (UP_4dA),
 P^1 -(2'-deoxyinosine 5'-)- P^4 -(uridine 5'-)tetraphosphate (UP_4dI) and
 P^1 -(2'-deoxycytidine 5'-)- P^4 -(uridine 5'-)tetraphosphate (UP_4dC).

THIS PAGE BLANK (USPTO)



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07H 21/00, A61K 31/70	A3	(11) International Publication Number: WO 98/34942 (43) International Publication Date: 13 August 1998 (13.08.98)
(21) International Application Number: PCT/US98/02702 (22) International Filing Date: 6 February 1998 (06.02.98) (30) Priority Data: 08/797,472 6 February 1997 (06.02.97) US 08/798,508 10 February 1997 (10.02.97) US (63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Applications US 08/797,472 (CIP) Filed on 6 February 1997 (06.02.97) US 08/798,508 (CIP) Filed on 10 February 1997 (10.02.97) (71) Applicant (for all designated States except US): INSPIRE PHARMACEUTICALS, INC. [US/US]; 4222 Emperor Boulevard #470, Durham, NC 27703 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): PENDERGAST, William [US/US]; 5815 Williamsburg Way, Durham, NC 27713 (US). YERKA, Benjamin, R. [US/US]; 1330 Hathaway Road, Raleigh, NC 27608 (US). RIDEOUT, Janet, L. [GB/US]; 3101 Morningside Drive, Raleigh, NC 27607		(US). SIDDIQI, Suhaib, M. [AT/US]; 1005 Paddock Drive, Raleigh, NC 27609 (US). (74) Agent: HALLUIN, Albert, P.; Howrey & Simon, 1299 Pennsylvania Avenue, N.W., Box 34, Washington, DC 20004-2402 (US). (81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> (88) Date of publication of the international search report: 6 January 2000 (06.01.00)
(54) Title: DINUCLEOTIDES AND THEIR USE (57) Abstract <p>The present invention relates to certain novel dinucleotides and formulations thereof which are highly selective agonists of the P2Y₂ and/or P2Y₄ purinergic receptor. They are useful in the treatment of chronic obstructive pulmonary diseases such as chronic bronchitis, PCD, cystic fibrosis, as well as prevention of pneumonia due to immobility. Furthermore, because of their general ability to clear retained mucus secretions and stimulate ciliary beat frequency, the compounds of the present invention are also useful in the treatment of sinusitis, otitis media and nasolacrimal duct obstruction. They are also useful for treatment of dry eye disease and retinal detachment.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

INTERNATIONAL SEARCH REPORT

Internati Application No
PCT/US 98/02702

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07H21/00 A61K31/70

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07H A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 96 40059 A (THE UNIVERSITY OF NORTH CAROLINA AT CHAPEL HILL) 19 December 1996 (1996-12-19) the whole document	1,3,6-8, 15-21
X	--- KANAVARIOTI A ET AL: "UNEXPECTEDLY FACILE SYNTHESIS OF SYMMETRICAL P1,P2-DINUCLEOSIDE-5' PYROPHOSPHATES" TETRAHEDRON LETTERS, vol. 32, no. 43, 21 October 1991 (1991-10-21), pages 6065-6068, XP002057243 cited in the application see page 6066, table 1 N = C or U --- -/-	1-3,6-8



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

22 February 1999

Date of mailing of the international search report

16. 03. 1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

SCOTT, J

INTERNATIONAL SEARCH REPORT

Internal: Application No
PCT/US 98/02702

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	CHEMICAL ABSTRACTS, vol. 125, no. 19, 4 November 1996 (1996-11-04) Columbus, Ohio, US; abstract no. 241071d, P.Z.DING ET AL.: "Oligomerization of Uridine Phosphorimidazolidones on Montmorillonite : a Model for the Prebiotic Synthesis of RNA on Minerals." page 435; column 2; XP002073884 abstract & ORIGINS LIFE EVOL. BIOSPHERE, vol. 26, no. 2, 1996, pages 151-171, ---	1,3,6-8
X	M-E.THEOCLITOU ET AL.: "Characterization of stress protein LysU. Enzymic synthesis of diadenosine 5',5'''-P1,P4-tetraphosphate (Ap4A) analogs by LysU." JOURNAL OF THE CHEMICAL SOCIETY, PERKIN TRANSACTIONS 1., no. 16, 21 August 1996 (1996-08-21), pages 2009-2019, XP002092434 LETCWORTH GB see page 2013, compounds 27,28,42,44 see page 2013, compounds 23-25,30-32 ---	1-3,6,8, 15
X	A.GURANOWSKI ET AL.: "Methylene and Halomethylene Analogues of Diadenosine 5',5'''-P1,P3-Triphosphate (ApppA) As Substrates or Inhibitors of ApppA-Degrading Enzymes." NUCLEOSIDES AND NUCLEOTIDES, vol. 14, no. 3-5, 1995, pages 731-734, XP002092438 See page 733, ApCH2ppA, ApCF2ppA ---	1-3,6,8, 15
X	WO 96 02554 A (GRUPPO LEPETIT S.P.A.) 1 February 1996 (1996-02-01) claims 1,4,8,14 ---	1,6-8,15
X	G.M.BLACKBURN ET AL.: "Novel Phosphonate and Thiophosphate Analogues of AP3A, Diadenosine 5',5'''-P1,P3-Triphosphate." TETRAHEDRON LETTERS., vol. 31, no. 39, 17 September 1990 (1990-09-17), pages 5637-40, XP002092436 OXFORD GB page 5637; examples 2A,2C ---	1,2,6,8, 15
	-/--	

INTERNATIONAL SEARCH REPORT

Internal Application No
PCT/US 98/02702

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	J.VISSCHER ET AL.: "Selective Cleavage of Pyrophosphate Linkages." NUCLEIC ACIDS RESEARCH., vol. 20, no. 21, 11 November 1992 (1992-11-11), pages 5749-5752, XP002073879 OXFORD GB page 5750, column 1, line 11 ---	1,6-8
X	E.HOLLER ET AL.: "Circular Dichroism and Ordered Structure of Bisnucleoside Oligophosphates and Their Zn ²⁺ and Mg ²⁺ Complexes." BIOCHEMISTRY, vol. 22, 1983, pages 4924-4933, XP002092437 See page 4924, compounds Gp5G, Gp6G, Gp3A, Gp3C, Gp3U ---	1-3,6,8
X	R.M.ORR ET AL.: "Inhibition of Human Leukaemic Thymidylate Kinase and L1210 Ribonucleotide Reductase By Dinucleotides of Adenosine and Thymidine and Their Phosphonate Analogues." BIOCHEMICAL PHARMACOLOGY, vol. 37, no. 4, 1988, pages 673-677, XP002092439 see the abstract, Ap2cpT, Ap3cpT, Ap4cpT, Ap3T, Ap4T, Ap5T, Ap6T ---	1,3,6,8
X	K.ONO ET AL.: "Inhibitory Effects of Diadenosine Polyphosphates and Diguanosine Polyphosphates on Terminal Deoxynucleotidyltransferase from Calf Thymus." BIOMEDICINE, vol. 36, no. 10, 1982, pages 414-419, XP002092440 See page 417, Gp5G and Gp6G ---	1,6,8
X	P.PLATEAU ET AL.: "Catabolism of Bis(5'-Nucleosidyl) Oligophosphates in Escherichia Coli : Metal Requirements and Substrate Specificity of Homogeneous Diadenosine-5',5''-P1,P4-tetraphosphate Pyrophosphohydrolase." BIOCHEMISTRY, vol. 24, 1985, pages 914-922, XP002092441 see abstract, compounds Ap3G, Ap3C, Gp3C ---	1-3,6,8
	-/--	

INTERNATIONAL SEARCH REPORT

Internal : Application No

PCT/US 98/02702

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	A.J.CHAVAN ET AL.: "Identification of N-Terminus Peptide of Human Granulocyte/Macrophage Colony Stimulating Factor as the Site of Nucleotide Interaction." BIOCHEMICAL AND BIOPHYSICAL RESEARCH COMMUNICATIONS, vol. 208, no. 1, 8 March 1995 (1995-03-08), pages 390-396, XP002092442 see p.391 8N3Ap4A	1-3,6,8
X	--- E.HAGEMEIER ET AL.: "High-Performance Liquid Chromatographic Method for Separation of Dinucleotides." JOURNNAL OF CHROMATOGRAPHY, vol. 237, no. 1, 1982, pages 174-177, XP002092443 see page 176, compounds Gp3A Ip3G, Ip3A, m7Gp3A, Ip5I	1-3,6,8
X	--- K.SCHEFFZEK: "Crystal Structure of the Complex of UMP/CMP Kinase from Dictyostelium Discoideum and the Bisubstrate Inhibitor P1-(5'-Adenosyl) P5-(5'-Uridyl) Pentaphosphate (UP5A) and Mg2+ at 2.2 Å : Implications for Water-Mediated Specificity." BIOCHEMISTRY, vol. 35, no. 30, 30 July 1996 (1996-07-30), pages 9716-9727, XP002092444 see abstract Up5A	1-3,6,8
X	--- S.STRIDH ET AL.: "Functional Analysis of Influenza RNA Polymerase Activity by the Use of Caps, Oligonucleotides and Polynucleotides." ANTIVIRAL RESEARCH, vol. 1, no. 2, 1981, pages 97-105, XP002092446 see page 99, table 1	1-3,6,8
X	--- J.PINTOR ET AL.: "Diinosine Polyphosphates, a Group of Dinucleotides with Antagonistic Effects on Diadenosine Polyphosphate Receptor." MOLECULAR PHARMACOLOGY, vol. 51, no. 2, February 1997 (1997-02), pages 277-284, XP002092447 see page 278, column 2, line 1, Ip3I, Ip4I, Ip5I --- -/--	1-3,6,8

INTERNATIONAL SEARCH REPORT

Internat. Application No

PCT/US 98/02702

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	CHEMICAL ABSTRACTS, vol. 110, no. 17, 24 April 1989 (1989-04-24) Columbus, Ohio, US; abstract no. 154770c, N.B.TARUSOVA ET AL.: "Organophosphorus Analogues of Biologically Active Compounds. XVI. Comparison of the Optical Properties of Zinc II Complexes of P1,P4-Bis(5'-Adenosyl)tetrphosphate and its Phosphonate Analogs." page 770; column 1; XP002092449 abstract & ZH. ORG. KHIM., vol. 24, no. 7, 1988, page 1474-80 ---	1-3,6,8
X	T.HATA ET AL.: "The Synthesis of alpha,gamma-Dinucleoside Triphosphates : The Confronted Nucleotide Structure Found at the 5'-Terminus of Eukaryote Messenger RNA." CHEMISTRY LETTERS, 1976, pages 987-990, XP002093657 the whole document ---	1-3,6,8
X	CHEMICAL ABSTRACTS, vol. 109, no. 1, 4 July 1988 (1988-07-04) Columbus, Ohio, US; abstract no. 6867d, Y.V.TUMANOV ET AL.: "Chemical Synthesis of Nucleoside-5'-Polyphospho-5'-Nucleosides." page 655; column 1; XP002092450 abstract & BIOORG. KHIM., vol. 13, no. 7, 1987, pages 921-927, ---	1-3,6,8
X	G.F.HUHN ET AL.: "Purification of Nucleoside-5'-diphosphates : A New Ion-Exchange Method." SEPARATION AND SCIENCE TECHNOLOGY, vol. 28, no. 11 & 12, 1993, pages 1959-1970, XP002093658 the whole document ---	1-3,6,8
A	"US ORPHAN STATUS FOR INSPIRE'S UTP" SCRIP, no. 2151, 2 August 1996 (1996-08-02), page 14 XP002032361 the whole document ---	1,15-21

	-/--	

INTERNATIONAL SEARCH REPORT

Internat Application No PCT/US 98/02702

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>LETHEM M I ET AL: "NUCLEOTIDE REGULATION OF GOBLET CELLS IN HUMAN AIRWAY EPITHELIAL EXPLANTS: NORMAL EXOCYTOSIS IN CYSTIC FIBROSIS"</p> <p>AMERICAN JOURNAL OF RESPIRATORY CELL AND MOLECULAR BIOLOGY, 55757, 20, vol. 9, 1993, pages 315-322, XP002055755 cited in the application the whole document</p> <p style="text-align: center;">---</p>	1,15-21
A	<p>US 5 292 498 A (BOUCHER JR RICHARD C) 8 March 1994 (1994-03-08) abstract</p> <p style="text-align: center;">---</p>	1,15-21
A	<p>OLIVIER K N ET AL: "ACUTE SAFETY AND EFFECTS ON MUCOCILIARY CLEARANCE OF AEROSOLIZED URIDINE 5-TRIPHOSPHATE + AND - AMILORIDE IN NORMAL HUMAN ADULTS"</p> <p>AMERICAN JOURNAL OF RESPIRATORY AND CRITICAL CARE MEDICINE, vol. 154, no. 1, July 1996 (1996-07), pages 217-223, XP002056055 cited in the application the whole document</p> <p style="text-align: center;">---</p>	1,15-21
A	<p>J.O.KLEIN: "Otitis Media."</p> <p>CLINICAL INFECTIOUS DISEASES, vol. 19, no. 5, November 1994 (1994-11), pages 823-833, XP002073880 cited in the application the whole document</p> <p style="text-align: center;">---</p>	1,15-21
A	<p>NUUTINEN J: "ACTIVATION OF THE IMPAIRED NASAL MUCOCILIARY TRANSPORT IN CHILDREN: PRELIMINARY REPORT"</p> <p>INTERNATIONAL JOURNAL OF PEDIATRIC OTORHINOLARYNGOLOGY, vol. 10, no. 1, October 1985 (1985-10), pages 47-52, XP000654070 the whole document</p> <p style="text-align: center;">---</p>	1,15-21
A	<p>BROWN H A ET AL: "EVIDENCE THAT UTP AND ATP REGULATE PHOSPHOLIPASE C THROUGH A COMMON EXTRACELLULAR 5'-NUCLEOTIDE RECEPTOR IN HUMAN AIRWAY EPITHELIAL CELLS"</p> <p>MOLECULAR PHARMACOLOGY, vol. 40, no. 5, November 1991 (1991-11), pages 648-655, XP002055754 cited in the application the whole document</p> <p style="text-align: center;">---</p> <p style="text-align: center;">-/--</p>	1,15-21

INTERNATIONAL SEARCH REPORT

Internat I Application No

PCT/US 98/02702

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>STEPINSKI J ET AL: "SYNTHESIS AND PROPERTIES OF P1,P2-, P1,P3- AND P1,P4-DINUCLEOSIDE DI-, TRI-AND TETRAPHOSPHATE MRNA 5'-CAP ANALOGUES" NUCLEOSIDES & NUCLEOTIDES, vol. 14, no. 3/05, 1995, pages 717-721, XP000608112 cited in the application see page 718, compounds 1-3</p>	1
A	<p>A.ZATORSKI ET AL.: "Chemical Synthesis of Benzamide Adenine Dinucleoide : Inhibition of Inosine Monophosphate Dehydrogenase (Types I and II)." JOURNAL OF MEDICINAL CHEMISTRY., vol. 39, 1996, pages 2422-2426, XP002073881 WASHINGTON US cited in the application the whole document</p>	1
A	<p>M.A.G.SILLERO ET AL.: "Dinucleosidetriphosphaase from Rat Liver." EUROPEAN JOURNAL OF BIOCHEMISTRY, vol. 76, 1977, pages 331-337, XP002073882 cited in the application the whole document</p>	1
A	<p>KAM-MUI EVA NG ET AL.: "The Action of a Water-Soluble Carbodiimide on Adenosine-5'-Polyphosphates." NUCLEIC ACIDS RESEARCH., vol. 15, no. 8, 1987, pages 3573-3580, XP002073883 OXFORD GB cited in the application the whole document</p>	1
A	<p>G.M.BLAKBURN ET AL.: "Synthesis, Physical, Chemical, and enzyme Studies on Bis-2,6-Diaminopurine B-D_Ribofuranoside P1,P4-Tetraphosphate." NUCLEOSIDES AND NUCLEOTIDES, vol. 10, no. 1-3, 1991, pages 549-551, XP002092448 the whole document</p>	1
P,X	<p>WO 98 03182 A (INSPIRE PHARMACEUTICALS INC ;JACOBUS KARLA M (US); LEIGHTON H JEFF) 29 January 1998 (1998-01-29) the whole document</p>	1-22

-/--

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 98/02702

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	WO 98 03177 A (INSPIRE PHARMACEUTICALS INC ;DRUTZ DAVID J (US); JABOBUS KARLA (US) 29 January 1998 (1998-01-29) the whole document ---	1-22
E	WO 98 15563 A (CODON PHARMACEUTICALS, INC.) 16 April 1998 (1998-04-16) the whole document -----	1-22

INTERNATIONAL SEARCH REPORT

Inter. Application No.

PCT/US 98/ 02702

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 16-21
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claim(s) 16-21
is(are) directed to a method of treatment of the human/animal
body, the search has been carried out and based on the alleged
effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such
an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

See further information sheet

1. ☒ As all required additional search fees were timely paid by the applicant, this International Search Report covers all
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☒ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

1. Claims: 1-4,6-8,15-21 partially

Compounds of formula I, where $X = O$, and $n+m = 0$; and uses thereof.

2. Claims: 1-4,6-8,15-21 partially ;

Compounds of formula I, where $X = O$, and $n+m = 1$; and uses thereof.

3. Claims: 1-4,6-8,15-22 partially ; 12,14 completely

Compounds of formula I, where $X = O$, and $n+m = 2$; and uses thereof.

4. Claims: 1-8,15-21 partially ; 13 completely

Compounds of formula I, where $X = O$, and $n+m = 3$; and uses thereof.

5. Claims: 1-8,15-22 partially

Compounds of formula I, where $X = O$, and $n+m = 4$; and uses thereof.

6. Claims: 1-4,6-8,15-21 partially ; 9 completely

Compounds of formula I, where $X = CH_2$; and uses thereof.

7. Claims: 1-4,6-8,15-21 partially ; 10 completely

Compounds of formula I, where $X = CF_2$; and uses thereof.

8. Claims: 1-4,6-8,15-21 partially ; 11 completely

Compounds of formula I, where $X = \text{imide}$; and uses thereof.

INTERNATIONAL SEARCH REPORT

Information on patent family members

Internal Application No

PCT/US 98/02702

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9640059 A	19-12-1996	US 5635160 A	03-06-1997
		AU 696845 B	17-09-1998
		AU 6176496 A	30-12-1996
		BR 9609063 A	26-01-1999
		CA 2223894 A	19-12-1996
		EP 0831777 A	01-04-1998
		JP 11506790 T	15-06-1999
		NO 975632 A	03-02-1998
		US 5935555 A	10-08-1999
WO 9602554 A	01-02-1996	AT 177431 T	15-03-1999
		AU 688552 B	12-03-1998
		AU 3076695 A	16-02-1996
		CA 2195091 A	01-02-1996
		DE 69508255 D	15-04-1999
		DE 69508255 T	14-10-1999
		EP 0773951 A	21-05-1997
		ES 2129893 T	16-06-1999
		FI 970081 A	08-01-1997
		HU 76331 A	28-08-1997
		JP 10502655 T	10-03-1998
		NO 965483 A	28-01-1997
		NZ 290127 A	27-05-1998
		ZA 9505826 A	21-02-1996
US 5292498 A	08-03-1994	WO 9408593 A	28-04-1994
		AU 2872592 A	09-05-1994
		AU 688258 B	12-03-1998
		EP 0663830 A	26-07-1995
		JP 8502078 T	05-03-1996
		NO 951340 A	02-06-1995
WO 9803182 A	29-01-1998	US 5763447 A	09-06-1998
		AU 3737597 A	10-02-1998
		CA 2261674 A	29-01-1998
		CN 1226169 A	18-08-1999
		EP 0938322 A	01-09-1999
		ZA 9706538 A	09-07-1998
WO 9803177 A	29-01-1998	US 5789391 A	04-08-1998
		AU 4408997 A	10-02-1998
		CA 2259552 A	29-01-1998
		CN 1227491 A	01-09-1999
		EP 0939637 A	08-09-1999
		US 5958897 A	28-09-1999
		US 5972904 A	26-10-1999
WO 9815563 A	16-04-1998	AU 4815197 A	05-05-1998
		EP 0934325 A	11-08-1999

THIS PAGE BLANK (USPTO)